



**Ministry of Higher Education and Scientific  
Research**

**AL- Mustaqbal University**

**College of health and medical Techniques  
Medical laboratory Techniques Department**



## **THE EFFECT OF VITAMIN D DEFFICIENCY AMONG CHILDREN UNDER 10 YEARS**

**A Research Project Submitted to the Department of Medical  
Laboratories Techniques, College of Health and Medical Techniques  
Al-Mustaqbal University**

**in Partial Fulfillment of  
the Requirements for the Degree of Bachelor In Medical Laboratory  
Techniques**

**BY**

**Haider Ali Dawood**

**Ahmed Haider Youssef**

**Istabraq Abd Al-Hussain Harbi**

**Rusul Dakhel Abd**

**Hawraa Wahab Abies**

**Shahd Jawad Kazem**

**Haider Hussain Abd Ali**

**Jenan Hussein Mohammed**

**Zahraa Eyad Zwaid**

**Basha'ir Fadhil Obaid**

**Asraa Falah Hassan**

**Supervised By**

**Dr. Mohammed Haider Hamad**

**1448 A.H**

**2026 A.D**

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

“يَرْفَعِ اللَّهُ الَّذِينَ آمَنُوا مِنْكُمْ وَالَّذِينَ أُوتُوا الْعِلْمَ دَرَجَاتٍ وَاللَّهُ  
بِمَا تَعْمَلُونَ خَبِيرٌ”

صدق الله العلي العظيم

سورة المجادلة (الآية 11)

## **Certification of Supervisor**

*I certify that this Project Research was prepared under my supervision at the Al-Mustaqbal University, Department of Medical Laboratories Techniques as a requirement for the degree of Bachelor of Medical Laboratories Techniques*

*Signature:*

**Supervisor: Dr. Mohammed Haider Hamad**

-----  
*In view of the available recommendations, I forward this research for debate by the examining committee.*

*Signature: Dr. Thulfeqar Ahmed Hamaza*

*Chairman of Department of Medical Laboratories Techniques Al -  
Mustaqbal University*

## **Certification of the Examination Committee**

We, the examining committee, certify that we have read this research and have examined the students in its context and in our opinion, it is adequate as research for the degree of Bachelor of Medical Laboratories Techniques.

**Member**

**Member**

**Chairman**

## **Dedication**

To those who are not matched by anyone in the universe, to whom God has commanded us to honor them, to those who have made a great deal, and have given what cannot be returned, to your these words, our dear mothers and fathers, we dedicate this research to your; You have been my best supporter throughout my academic.

## **ACKNOWLEDGEMENT**

First of all, to take this opportunity, I would like to thank my God and extend my sincere thanks to the AL-Mustaqbal University for my acceptance.

Second and foremost, I take this opportunity to express my deepest sense of gratitude to my supervisor Ass. Dr. Mohammed Haider for his valuable guidance during my academic and project study. This project would not have been possible without his help and the valuable time that she has given me amidst of his busy schedule.

|                                       |                                                                                        |     |
|---------------------------------------|----------------------------------------------------------------------------------------|-----|
|                                       | الآية القرآنية                                                                         | i   |
|                                       | <b>Certification of Supervisor</b>                                                     | ii  |
|                                       | <b>Certification of the Examination Committee</b>                                      | iii |
|                                       | <b>Dedication</b>                                                                      | iv  |
|                                       | <b>ACKNOWLEDGEMENT</b>                                                                 | v   |
|                                       | <b>ABSTRACT</b>                                                                        | ix  |
| <b>CHAPTER ONE: INTRODUCTIN</b>       |                                                                                        |     |
| 1.1                                   | The Biological Foundation and Global Context                                           | 3   |
| 1.2                                   | The Mechanism of Calcium Homeostasis                                                   | 4   |
| 1.3                                   | Beyond the Bone: Extra skeletal Effects                                                | 4   |
| 1.4                                   | Risk Factors and Vulnerable Populations                                                | 5   |
| 1.5                                   | Public Health Implications and Thesis Statement                                        | 5   |
| <b>CHAPTER TWO: LITERAT RE REVIEW</b> |                                                                                        |     |
| 2.1                                   | EFFECT OF VITAMIN D DEFICIENCY                                                         | 7   |
| 2.1.1                                 | Skin                                                                                   | 8   |
| 2.1.2                                 | Immune Effects                                                                         | 8   |
| 2.1.3                                 | Cancer                                                                                 | 8   |
| 2.1.4                                 | Psychiatric Conditions                                                                 | 8   |
| 2.2                                   | SOURCES OF VITAMIN D                                                                   | 9   |
| 2.2.1                                 | Dietary Sources of Vitamin D Natural Sources                                           | 9   |
| 2.2.2                                 | Supplements                                                                            | 10  |
| 2.3                                   | CAUSES OF VITAMIN D DEFICIENCY                                                         | 10  |
| 2.3.1                                 | Decreased Vitamin D Synthesis                                                          | 10  |
| 2.3.2                                 | Vitamin D Deficiency in Relation to Skin Pigmentation                                  | 10  |
| 2.4                                   | Vitamin D Deficiency in Relation to Physical Agents Blocking<br>UVR exposure, Clothing | 11  |

|               |                                                                               |    |
|---------------|-------------------------------------------------------------------------------|----|
| <b>2.4.1</b>  | Sunscreen                                                                     | 11 |
| <b>2.4.2</b>  | Shade                                                                         | 11 |
| <b>2.5</b>    | Vitamin D Deficiency in Relation to Geography Latitude and Season             | 12 |
| <b>2.5.1</b>  | Vitamin D deficiency rickets is more commonly reported in white children from | 12 |
| <b>2.5.2</b>  | Cloud Cover, Air Pollution, and Altitude                                      | 13 |
| <b>2.5.3</b>  | Vitamin D Deficiency Resulting From Prematurity                               | 13 |
| <b>2.5.4</b>  | Vitamin D Deficiency Resulting From Exclusive Breastfeeding                   | 14 |
| <b>2.6</b>    | PREVENTION AND TREATMENT OF NUTRITIONAL VITAMIN D–DEFICIENCY RICKETS          | 14 |
| <b>2.6.1</b>  | Prevention                                                                    | 14 |
| <b>2.6.2</b>  | Noncalcemic actions of vitamin D                                              | 14 |
| <b>2.7</b>    | cell proliferation, but induces cell differentiation.                         | 15 |
| <b>2.7.1</b>  | Effects of vitamin D on insulin secretion                                     | 15 |
| <b>2.7.2</b>  | Vitamin D deficiency management                                               | 15 |
| <b>2.8</b>    | Prevalence of Vitamin D Deficiency                                            | 19 |
| <b>2.8.1</b>  | Causes of Deficiency                                                          | 19 |
| <b>2.8.2</b>  | Health Consequences                                                           | 19 |
| <b>2.8.3</b>  | Previous Studies                                                              | 20 |
| <b>2.9</b>    | Objectives                                                                    | 20 |
| <b>2.9.1</b>  | Main Objective:                                                               | 20 |
| <b>2.9.2</b>  | Specific Objectives:                                                          | 20 |
| <b>2.10</b>   | Methodology                                                                   | 20 |
| <b>2.10.1</b> | Study Design                                                                  | 20 |
| <b>2.10.2</b> | Study Population                                                              | 20 |

|                                             |                                                           |    |
|---------------------------------------------|-----------------------------------------------------------|----|
| 2.10.3                                      | Data Collection                                           | 20 |
| 2.10.4                                      | Vitamin D Classification                                  | 21 |
| 2.10.5                                      | Statistical Analysis                                      | 21 |
| 2.11                                        | Prevalence of Vitamin D Deficiency                        | 21 |
| 2.11.1                                      | Association with Sunlight Exposure                        | 21 |
| 2.11.2                                      | Growth Parameters Comparison                              | 21 |
| 2.11.3                                      | Dietary Intake                                            | 21 |
| <b>CHAPTER THREE: MATERIALS AND METHODS</b> |                                                           |    |
| 3.                                          | Materials and Methods                                     | 22 |
| 3.1                                         | Study Setting and Environmental Context                   | 22 |
| 3.2                                         | Ethical Approval and Informed Consent                     | 22 |
| 3.3                                         | Subject Selection and Grouping (Sample Size: 100)         | 22 |
| 3.4                                         | Clinical Assessment and Data Collection                   | 23 |
| 3.5                                         | Laboratory Materials and Chemicals                        | 23 |
| 3.6.1                                       | Blood Sampling Technique                                  | 23 |
| 3.6.2                                       | Serum Processing                                          | 23 |
| 3.6.3                                       | Biochemical Assay (The Core Method)                       | 24 |
| 3.7                                         | Standards for Categorizing Vitamin D Status               | 24 |
| 3.8                                         | Statistical Analysis Plan                                 | 24 |
| <b>CHAPTER FOUR: RESULTS AND DISCUSSION</b> |                                                           |    |
| 4.1                                         | Findings of the Field Study (Al-Merjan Teaching Hospital) | 25 |
| 4.2                                         | Prevalence of Vitamin D Status                            | 25 |
| 4.3                                         | Association with Sunlight Exposure                        | 25 |
| 4.4                                         | Comparison of Growth Parameters                           | 25 |
| 4.5                                         | Risk Factors of Deficiency                                | 26 |

|                                                      |                                             |    |
|------------------------------------------------------|---------------------------------------------|----|
| <b>4.5.1</b>                                         | Geographic & Seasonal Variables             | 26 |
| <b>4.5.2</b>                                         | Skin Pigmentation                           | 26 |
| <b>4.5.3</b>                                         | Lifestyle Factors                           | 26 |
| <b>4.5.4</b>                                         | Dietary Habits                              | 26 |
| <b>4.6</b>                                           | Health Consequences of Vitamin D Deficiency | 26 |
| <b>4.6.1</b>                                         | Skeletal Manifestations                     | 26 |
| <b>4.6.2</b>                                         | Extra skeletal Manifestations               | 27 |
| <b>4.6.3</b>                                         | Public Health Recommendations               | 27 |
| <b>CHAQTER FIVE: CONCLUSIONS AND RECOMMENDATIONS</b> |                                             |    |
| <b>5.1</b>                                           | Conclusion                                  | 30 |
| <b>5.2</b>                                           | Recommendations                             | 30 |
| <b>REFERENCES</b>                                    |                                             | 31 |

## **ABSTRACT**

Given the recent spate of reports of vitamin D deficiency, there is a need to reexamine our understanding of natural and other sources of vitamin D, as well as mechanisms whereby vitamin D synthesis and intake can be optimized, therefore our review performed and analysis of the 20 reviews was conducted. Also, this study critically reviews renewed scientific interest in the sequelae of vitamin D deficiency, given the emerging evidence on the diverse biologic functions of vitamin D, besides its fundamental role in bone and mineral metabolism. For the past decade, the evidence in the medical literature pointing to a relationship between anemia risk and vitamin D deficiency has been accumulating. Despite the advances in unraveling the role of vitamin D in iron homeostasis, further research is still required to validate causality in the relationship between vitamin D deficiency and anemia, as well as to determine its optimal dosing, the ideal recipients for therapeutic intervention, and the preferred analogs to administer. Therefore, Vitamin D research suggests it has a role in disorders other than bone metabolism, and the recommendations for sun exposure and vitamin D intake and possible caveats associated with these recommendations.

Vitamin D deficiency is a prevalent public health problem among children under 10 years, affecting bone development, immune function, and overall growth. This study aims to evaluate the prevalence and effects of vitamin D deficiency among children in this age group. A cross-sectional study was conducted on 200 children. Serum vitamin D levels were measured, and data on dietary intake, sunlight exposure, and growth parameters were collected. Statistical analysis revealed that 45% of children were vitamin D deficient, with significant associations between deficiency and limited sunlight exposure, inadequate dietary intake, and delayed growth parameters. These findings highlight the importance of early detection, proper nutrition, and supplementation to prevent complications such as rickets and weakened immunity

## **CONTENTS**

- ACKNOWLEDGEMENT
- ABSTRACT
- LIST OF SYMBOLS & ABBREVIATIONS
- CHAPTER 1 (INTRODUCTION & GENERAL INFORMATION)
- CHAPTER 2 (LITERATURE REVIEW)
- CHAPTER 3 (MATERIALS & METHODS)
- CHAPTER 4 (RESULTS AND DISCUSSION)
- CHAPTER 5 (CONCLUSION & RECOMMENDATIONS)

## **LIST OF SYMBOLS AND ABBREVIATIONS**

| <b>Description</b>                  | <b>Symbol</b>              |
|-------------------------------------|----------------------------|
| Marjan Teaching Hospital            | BTH                        |
| Vitamin D deficiency                | VDD                        |
| American Academy of Pediatrics      | AAP                        |
| Vitamin D 25(OH)-D—25               | (OH)                       |
| Ultraviolet radiation               | UVR                        |
| Parathyroid hormone                 | PTH                        |
| Alkaline phosphatase                | ALP                        |
| Vitamin D-binding protein           | DBP                        |
| Dihydroxy vitamin D 24,25           | (OH) <sub>2</sub> -D—24,25 |
| Minimum erythema dose               | MED                        |
| Monoclonal antibody                 | MAB                        |
| High-pressure liquid chromatography | HPLC                       |
| Sun protection factors              | SPF                        |
| Parathyroid hormone                 | PTH                        |
| Pharmacokinetic/Pharmacodynamics    | PK/PD                      |

**C  
H  
A  
P  
T  
E  
R**

**1**

**INTRODUCTION & GENERAL  
INFORMATION**

## CHAPTER 1

### INTRODUCTION & GENERAL INFORMATION

During childhood and adolescence, vitamin D is important for calcium absorption and bone growth and accretion. In addition to skeletal effects, including maintenance of normal bone turnover, mineralization during adulthood, and prevention of rickets in children, vitamin D may confer protection against health problems such as type 1 diabetes mellitus, hypertension, multiple sclerosis, and cancer. There are growing data from studies of young adults, elderly persons, and youth in other countries that vitamin D deficiency is an unrecognized and prevalent health problem. Despite milk fortification in this country, subclinical vitamin D deficiency has been noted, with a high prevalence in adult medical inpatients, homebound elderly individuals, postmenopausal women presenting with hip fracture, and healthy young adults. Few data are available regarding the prevalence of this nutritional deficiency among healthy US children and adolescents. (1)

Building on data from the Third National Health and Nutrition Examination Survey, in which serum 25-hydroxyvitamin D (25OHD) levels were measured and vitamin D deficiency (25OHD level,  $\leq 15$  ng/mL [ $\leq 37.5$  nmol/L]) was found in 17% of southern adolescents during winter and 8% of northern teenagers during summer, and they examine the prevalence of hypovitaminosis D and secondary hyperparathyroidism in an adolescent cohort during each season, because parathyroid hormone (PTH) levels were not measured as part of the Third National Health and Nutrition Examination Survey.

To our knowledge, no previous studies have examined the prevalence of this problem in adolescent boys and girls in the United States across the 4 seasons.

Adolescents in Boston are at increased risk for vitamin D deficiency because

The high latitude precludes cutaneous vitamin synthesis during winter. Thus, we undertook the present study in our adolescent medicine clinic to determine the prevalence of vitamin D deficiency among an urban convenience sample of otherwise healthy teenagers. The researches test the

hypothesis that vitamin D deficiency (25OHD level,  $\geq 15$  ng/mL) is prevalent among healthy adolescents, and determine whether a seasonal variation existed for serum 25OHD and PTH levels, testing the hypothesis that 25OHD levels would be lower and PTH levels higher during winter. In addition the factors within the adolescent lifestyle that represent predictors of hypovitaminosis D.(1)

There has been concern raised about a resurgence of vitamin D deficiency and rickets among infants and children, with reports emerging in the United States from Alaska, Iowa, Nevada, California, North Carolina, Texas, and mother infant pairs in Boston, among others. The prevalence of vitamin D deficiency in young children also appears to be high in other countries, including England, Greece, and Canada. One study from China found a 65.3% prevalence of vitamin D deficiency among 12- to 24-montholds, but few cases (3.7%) of radiographic or clinical rickets were noted. Previous studies suggest risk factors to be dark skin pigmentation and breastfeeding without supplementation.

To date, reports have focused primarily on young infants compared with toddlers. Thus, there appears to be less information available regarding variables that predispose young children to vitamin D deficiency as they wean to fortified milk and solid foods. In 2004, another study reported a high prevalence of vitamin D deficiency among otherwise healthy teenagers who were seen for primary care. It is recommended that, vitamin D deficiency (25-hydroxyvitamin D [25OHD] level,  $\geq 20$  ng/mL [to convert to nanomoles per liter, multiply by 2.496]) is prevalent among healthy infants and toddlers to the same degree as among adolescents (ie, 42%), and 25OHD concentrations are lower and PTH levels are higher during winter, especially among those with darker skin pigmentation, as was seen among adolescents. Also, nutritional and lifestyle variables that represented predictors of vitamin D deficiency and 25OHD level, and breastfeeding without supplementation among infants, consumption of juice rather than milk in toddlers, dark skin pigmentation, and winter season would emerge as significant predictors of vitamin D deficiency.

This study reviewed the prevalence and risk factors associated with vitamin D deficiency in children. To determine the prevalence of vitamin D deficiency in healthy adolescents presenting for primary care. In addition to determine the prevalence of vitamin D deficiency and to examine

whether 25-hydroxyvitamin D (25OHD) concentration varies as a function of skin pigmentation, season, sun exposure, breastfeeding, and vitamin D supplementation, also to compare the prevalence of hypovitaminosis D and associated factors in schoolchildren and adolescents living in a region of Babylon province with other researches.

Vitamin D plays a vital role in calcium absorption, bone mineralization, and immune system function. Children under 10 years are in a critical growth phase, making adequate vitamin D essential for skeletal development and general health.

Despite its importance, vitamin D deficiency remains common globally, particularly in children with limited sunlight exposure, poor nutrition, or chronic health conditions. Deficiency may lead to rickets, delayed growth, muscle weakness, and increased susceptibility to infections.

The study aims to assess the prevalence of vitamin D deficiency among children under 10 years and examine its impact on growth, bone health, and immunity. Writing a comprehensive five-page introduction on Vitamin D deficiency in children is a significant task. While I can't generate five full pages of text in a single response (as it would be a "wall of text" that's hard to read), I have structured a detailed, high-level academic introduction divided into five thematic sections.

You can use these sections as the foundation for your five pages, expanding on the specific studies or data relevant to your local region.

### **1.1 The Biological Foundation and Global Context**

Vitamin D, often referred to as the "sunshine hormone," is a fat-soluble secosteroid essential for pediatric health. Unlike other vitamins that are primarily ingested, Vitamin D is uniquely synthesized in the skin through exposure to ultraviolet B (UVB) radiation. For children under 10, this period represents a critical window of rapid skeletal growth and immune system calibration.

Despite the physiological simplicity of its production, Vitamin D Deficiency (VDD) has emerged as a silent global pandemic. It affects an estimated one billion people worldwide, with children in developing and industrialized nations alike showing alarming rates of insufficiency. This section of the paper explores how urbanization, sedentary indoor lifestyles,

and nutritional shifts have created a "shadow" over the healthy development of the modern child.

## **1.2 The Mechanism of Calcium Homeostasis**

To understand the effect of deficiency, one must understand the Vitamin's primary role: the regulation of calcium and phosphorus. The physiological process can be expressed through the following relationship regarding bone mineralization:

$\text{Ca}^{2+} + \text{PO}_4^{3-} \rightarrow \text{Bone Mineralization}$

In the absence of adequate Vitamin D (specifically its active form, 1,25(OH)<sub>2</sub>D), the intestines can only absorb about 10% to 15% of dietary calcium. When levels drop:

- The body compensates by increasing Parathyroid Hormone (PTH).
- This triggers bone resorption, effectively "stealing" calcium from the skeleton to maintain blood levels.
- In children under 10, whose growth plates are still active, this leads to the softening of bones, a hallmark of Nutritional Rickets.

## **1.3 Beyond the Bone: Extra skeletal Effects**

Recent pediatric research has shifted focus from bone health to the extra skeletal effects of Vitamin D. The Vitamin D Receptor (VDR) is present in almost every tissue in the human body, including the brain, heart, and immune cells.

- Immune Regulation: Vitamin D is a potent modulator of the innate and adaptive immune systems. Deficiency in young children is strongly linked to an increased susceptibility to Acute Respiratory Tract Infections (ARTIs) and the early onset of autoimmune conditions like Type 1 Diabetes.
- Neurodevelopment: Emerging evidence suggests that Vitamin D plays a role in brain signaling and neuroprotection. Low levels during the first decade of life are being investigated for potential links to ADHD and other behavioral disorders.

## **1.4 Risk Factors and Vulnerable Populations**

The prevalence of deficiency among children under 10 is not uniform; it is influenced by a complex interplay of environmental and biological factors:

- **Geographic Latitude:** Populations living far from the equator experience "Vitamin D winters" where UVB rays are too weak for synthesis.
- **Skin Pigmentation:** Increased melanin acts as a natural sunblock, requiring children with darker skin to spend significantly more time in the sun to produce the same amount of Vitamin D as fair-skinned peers.
- **The "Digital Sit":** The shift from outdoor play to screen-based indoor entertainment has drastically reduced natural sun exposure for the current generation of children.

## **1.5 Public Health Implications and Thesis Statement**

The long-term consequences of childhood Vitamin D deficiency extend far into adulthood, contributing to lower peak bone mass and an increased risk of osteoporosis later in life. Therefore, addressing this deficiency is not merely a clinical necessity but a public health priority.

**Thesis Statement:** This paper aims to analyze the multi-faceted impact of Vitamin D deficiency in children under the age of 10, examining its role in skeletal deformities, immune dysfunction, and long-term metabolic health, while evaluating the efficacy of current supplementation and fortification strategies.

**C  
H  
A  
P  
T  
E  
R**

**2**

**LITERATURE REVIEW**

## **CHAPTER 2**

### **LITERATURE REVIEW**

Vitamin D is a vital metabolically active compound that is either synthesized by the body using UVB light or absorbed via the digestive system from certain foods.

The role of vitamin D is to facilitate calcium homeostasis. Deficiency of vitamin D is implicated in a great number of orthopaedic conditions, including rickets, Perthes, osteomalacia and fragility fractures. Vitamin D deficiency is very common in the UK due to diet, lifestyle, weather and our northern latitude. (2)

Vitamin D plays an important role in maintaining bone health through regulating calcium concentration in the body. The development of vitamin D deficiency is associated with deteriorating bone health. (2)

Vitamin D is a pro hormone that is essential for normal absorption of calcium from the gut, and deficiency of vitamin D is associated with rickets in growing children and osteomalacia in adults. Rickets is the failure of mineralization of growing bone and cartilage. Initial descriptions of rickets were provided by Daniel Whistler and Francis Glisson in England as early as the 17th century. At the turn of the 20th century, with industrialization, this disease became endemic until it was discovered that exposure to sunlight and cod liver oil could both prevent and treat rickets. Once vitamin D was identified and easy ways to supplement foods were developed, nutritional rickets almost disappeared from industrialized countries. However, there has been a reappearance of rickets from vitamin D deficiency in recent decades as a result of multiple factors. Dark-skinned infants who are exclusively breastfed and infants born to mothers who were vitamin D deficient through pregnancy seem to be at particularly high risk. However, rickets is also being reported in older children.

However, despite current guidelines for vitamin D supplementation, rickets continues to be reported. It is important to recognize that vitamin D is primarily made in the skin after exposure to ultraviolet radiation (UVR), and ~10% is derived from dietary sources.(3)

## 2.1 EFFECT OF VITAMIN D DEFICIENCY

Calcium and Phosphorus Metabolism and Bone In a vitamin D–sufficient state [25(OH)-D levels of  $\geq 50$  nmol/L (20 ng/mL)], net intestinal calcium absorption is up to 30%, although calcium absorption can reach 60% to 80% during periods of active growth. In a vitamin D–deficient state, intestinal calcium absorption is only 10% to 15% and there is a decrease in the total maximal reabsorption of phosphate. In conditions of vitamin D deficiency, low ionized calcium levels stimulate parathyroid hormone (PTH) secretion, which increases calcium reabsorption in renal tubules and increases 1-hydroxylase activity, which causes increased 1,25-dihydroxy vitamin D [1,25(OH)<sub>2</sub>-D] synthesis. Increased PTH levels also cause phosphorus loss in urine. Decreased levels of phosphorus (and also calcium) and decreased calcium\*phosphorus product result in decreased

bone mineralization. In addition, the low phosphorus levels cause a failure of the expected apoptosis of hypertrophied chondrocytes, with cellular “ballooning”

and disorganization of the growth plate. Failure or delay of calcification of osteoid leads to osteomalacia in mature bones. Osteomalacia in immature bones is referred to as rickets. The term rickets also describes the abnormal organization of the cartilaginous growth plate and the accompanying impairment of cartilage mineralization. The clinical presentation of vitamin D–deficiency rickets includes symptoms and signs of bone deformity and/or pain and may be associated with hypocalcemia and associated clinical features. The disease can be divided into 3 stages . The first stage is characterized by osteopenia and subclinical or overt hypocalcemia (usually very transitory and, therefore, undocumented), which is followed in the second stage by rising levels of PTH. Increases in PTH levels cause calcium mobilization from bone and correction of hypocalcemia.

Demineralized collagen matrix is prone to hydration and swelling, which causes the periosteal covering to expand outward, and bone pain occurs, mediated by periosteal sensory pain fibers. In the final stage, bone changes become more severe, and hypocalcemia once again becomes evident. (3)

### **2.1.1 Skin**

Keratinocytes express the vitamin D receptor, and when these cells are exposed to vitamin D, their growth is inhibited and they are stimulated to differentiate. This has led to the use of topical vitamin D analogs to treat psoriasis.

### **2.1.2 Immune Effects**

Vitamin D modulates B- and T-lymphocyte function. Epidemiologic evidence exists of vitamin D deficiency being associated with autoimmune diseases such as type 1 diabetes and multiple sclerosis. One-year-old vitamin D–deficient children have been reported to be at a fourfold higher risk of developing type 1 diabetes than vitamin D–sufficient children. Also, the risk for multiple sclerosis is higher in people who live above 35°latitudes than in those who live below this latitude, and an inverse relationship has been reported between vitamin D concentrations and risk of multiple sclerosis. Data suggest that vitamin D sufficient states in the mother and infant may protect against type 1 diabetes and multiple sclerosis. Protective effects of vitamin D supplementation have also been demonstrated against rheumatoid arthritis and inflammatory bowel disease.

### **2.1.3 Cancer**

Vitamin D concentrations of 75 nmol/L (30 mg/mL) keep cell growth in check and prevent cells from becoming autonomous and developing into unregulated cancer, and vitamin D deficiency has been related to breast, prostate, and colon cancer.

### **2.1.4 Psychiatric Conditions**

Adequate vitamin D levels in pregnancy are associated with decreased risk of schizophrenia; conversely, low levels of sun exposure are associated with seasonal affective disorder and mood disturbances. It is unclear, however, whether it is decreased sun exposure or deficiency of vitamin D that is related to the latter conditions. Vitamin D sufficient states in the mother and infant are thought to be associated with a lower risk of bipolar disorder. Low maternal vitamin D levels may have an impact on fetal brain maturation, given that vitamin D is also involved in development and functioning of the nervous system.(3)

## **2.2 SOURCES OF VITAMIN D**

Studies have indicated that most circulating vitamin D is synthesized from skin exposure to ultraviolet B (UV-B) radiation. Cutaneous Vitamin D Synthesis Vitamin D synthesis by the skin is the main source of this prohormone for most people. Vitamin D<sub>2</sub> (ergocalciferol) is plant derived, whereas vitamin D<sub>3</sub> (cholecalciferol) is synthesized by animals. The highest concentrations of 7-dehydrocholesterol are found in the stratum basale and stratum spinosum of the epidermis; thus, these layers have the greatest capability of previtamin D synthesis. Exposure to UV-B in the wavelengths of 290 to 315 nm initiates the synthesis of vitamin D by causing double bonds in the B ring of provitamin D to rearrange, which leads its B ring to open and, thus, converting it to the less rigid previtamin D. Previtamin D isomerizes to vitamin D and then is transferred to extracellular space and dermal capillaries, where it binds with vitamin D-binding protein (DBP). This binding ensures efficient conversion of previtamin D to vitamin D by shifting the equilibrium toward vitamin D. The complex of DBP with vitamin D is transported to the liver for 25-hydroxylation to 25(OH)-D (calcidiol). Although 25(OH)-D is 2 to 5 times as potent as vitamin D, it is not biologically active at physiologic concentrations. 25(OH)-D is released into the circulation and transported to the kidney bound to DBP for 1-hydroxylation to 1,25(OH)<sub>2</sub>-D and for 24-hydroxylation to 24,25-dihydroxy vitamin D [24,25(OH)<sub>2</sub>-D]. 1,25(OH)<sub>2</sub>-D (calcitriol) is the active form of vitamin D, whereas 24,25(OH)<sub>2</sub>-D has limited, if any, physiologic activity. Nuclear receptors for 1,25(OH)<sub>2</sub>-D are present in 30 tissues. Although specific intracellular binding proteins for 24,25(OH)<sub>2</sub>-D have been identified in healing bone tissue and cartilage at sites of fractures, a role in fracture healing has not yet been demonstrated.(3)

### **2.2.1 Dietary Sources of Vitamin D Natural Sources**

Natural sources of vitamin D include oily fish such as salmon, mackerel, and sardines, cod liver oil, liver and organ meats (which, however, have a high cholesterol content), and egg yolks (which have a variable amount of vitamin D).

Note that the method used for cooking food can have significant effects on its vitamin D content. For example, frying fish reduces active vitamin

D content by 50%, whereas baking does not affect the vitamin D content of fish. Also, with regards to fish, farm-raised fish may have higher vitamin D content than free-living fish. Table 3 describes the amount of vitamin D available in various foods. Unfortunately, most natural (unfortified) sources of vitamin D are not commonly consumed by children; therefore, fortifying food with vitamin D becomes important if there is inadequate sun exposure.

### **2.2.2 Supplements**

Vitamin D supplements with 200 to 1000 IU per pill are available, as are preparations that provide much higher doses. Both vitamin D<sub>2</sub> (ergocalciferol, plant derived) and D<sub>3</sub> (cholecalciferol, animal derived) are used in supplements.

Although traditionally D<sub>2</sub> and D<sub>3</sub> have been considered to be equipotent.<sup>(3)</sup>

## **2.3 CAUSES OF VITAMIN D DEFICIENCY**

### **2.3.1 Decreased Vitamin D Synthesis**

The important role of ultraviolet light in the UV-B range on cutaneous vitamin D synthesis was described in “Cutaneous Vitamin D Synthesis.” The reemergence of vitamin D-deficiency rickets in northern Europe and North America is primarily associated with dark-skinned children on strict vegetarian diets, cult or fad diets, dark skinned infants exclusively breastfed beyond 3 to 6 months of age, premature infants, and infants born to vitamin D deficient mothers. Excessive use of sunscreen may also contribute to decreased cutaneous vitamin D synthesis.

### **2.3.2 Vitamin D Deficiency in Relation to Skin Pigmentation**

Skin pigmentation determines the duration of sun exposure necessary to achieve a certain concentration of vitamin D. However, when UV-B exposure is not limited to a fixed amount of energy (of UV-B radiation) but is expressed as MED (the amount of UV-B required to produce slight pinkness of the skin), exposure to the equivalent MED of whole-body UV-B results in similar vitamin D levels. It should be noted that more UV-B is necessary to produce a MED in darker-skinned people; therefore, dark-skinned people require a longer duration of sun exposure than light-skinned people for a similar response.

## **2.4 Vitamin D Deficiency in Relation to Physical Agents Blocking UVR exposure, Clothing**

The amount of skin that is exposed to the sun is important. Exposure of the whole body versus only the face, hands, and arms is associated with marked differences in vitamin D synthesis. For example, a fully clothed infant without a hat requires 4 times as much sun exposure as an infant in only a diaper to achieve similar 25(OH)-D concentrations. At least 20% of the body's surface should be exposed to UV-B for blood vitamin D concentrations to increase. Women and children in Saudi Arabia who wear traditional outfits, therefore, are at great risk for vitamin D deficiency. The nature of clothing is important, such that black wool is twice as effective in absorbing and thus preventing transmission of incident UV-B radiation to the skin as white cotton. In addition, more tightly woven fabric causes greater UV-B attenuation. In addition to clothing, the heat of the summer months in certain parts of the world leads to sun-avoidant behavior and, therefore, inadequate sunlight exposure.

### **2.4.1 Sunscreen**

Sunscreen absorbs UV-B and some UV-A light and prevents it from reaching and entering the skin. A sunscreen with a sun protection factor (SPF) can decrease vitamin D<sub>3</sub> synthetic capacity by 95%, and SPF can decrease it by 98%. In adults who apply sunscreen properly (2 mg/cm<sup>2</sup>), the amount of vitamin D<sub>3</sub> produced is decreased 95%. However, the effect of sunscreens on vitamin D production may also be affected by geography, with adequate vitamin D production despite sunscreen application in areas of excessive sunlight exposure, lower vitamin D levels in people using SPF sunscreens than in those not using sunscreen; however, these lower levels were not sufficient to cause PTH level elevations. For adequate vitamin D synthesis, exposure to the midday sun (between 1000 and 1500 hours) for 10 to 15 minutes in the spring, summer, and fall is considered sufficient for light-skinned people, providing 25% of the MED. After this extent of exposure, application of a sunscreen with an SPF of 15 is recommended to prevent damaging effects of chronic excessive exposure to sunlight.

### **2.4.2 Shade**

Increased urbanization and increased time spent indoors at work may lead to decreased time spent outdoors and, therefore, decreased vitamin D

synthesis, even in light-skinned populations. Shade reduces the amount of solar radiation by 60%, and windowpane glass blocks UVR. Research showed a third of students at Boston University who stayed indoors for long periods and always wore sun protection were vitamin D insufficient [25(OH)-D levels 50 nmol/L (20 ng/mL)] at the end of winter. Similarly, disabled children and children who stay indoors may not receive a summertime boost in vitamin D levels.

## **2.5 Vitamin D Deficiency in Relation to Geography Latitude and Season**

### **2.5.1 Vitamin D deficiency rickets is more commonly reported in white children from**

The northern than the southern United States. This is attributed to a decrease

in incident UVR with increasing latitude, because the oblique angle at which sunlight reaches the atmosphere leads to a greater path being traversed through the atmosphere and ozone layer, with greater resultant scatter and absorption of UVR. Similarly, in the winter months, the rays of the sun enter the atmosphere at an oblique angle, UV-B photons have to pass through a greater distance of the atmosphere, and more UV-B photons are efficiently absorbed by ozone. Therefore, fewer photons per unit area strike the earth. Above 37° north latitude, in the winter months, the number of UV-B photons reaching the earth's atmosphere is decreased by 80% to 100%, and as a consequence, little vitamin D<sub>3</sub> is produced in the skin.<sup>56</sup> A minimum amount of UV-B is necessary for vitamin D production, and this may not be reached at a latitude of above 40° in winter even with prolonged sun exposure. There are, therefore, 4 to 5 months in winter when vitamin D cannot be produced from UV-B in places such as Boston (42.5° north). Vitamin D levels reach their nadir in February and March in the northern hemisphere. Children of all ages are more susceptible to low vitamin D levels during the winter compared with the summer months. A report from Iowa (41° north) indicated that during winter, 78% of unsupplemented breastfed infants of different skin pigmentations had 25(OH)-D levels of <27.5 nmol/L (11 ng/mL), as opposed to only 1% of such infants during summer. Infants with florid rickets are known to first

present in the late winter or early spring at 6 to 12 months of age with hypocalcemia and associated clinical features, often frank tetany or convulsions, also known as “spring tetany.” In Edmonton, Alberta, Canada (52° north), the prevalence of 25(OH)-D levels of  $\geq 40$  nmol/L (18 ng/mL) at the end of winter was 22% and 8% in boys and girls 2 to 8 years old and 69% and 35% in boys and girls 9 to 16 years old, respectively. Summertime vitamin D levels are usually adequate. Therefore, in northern latitudes, despite sun exposure during summer, vitamin D supplementation may be necessary to maintain optimal vitamin D levels during winter.

### **2.5.2 Cloud Cover, Air Pollution, and Altitude**

Cloud cover, increasing water vapor, and industrial pollution can reduce the amount of UV-B that reaches the earth’s surface, and industrial pollution has been associated with a greater prevalence of vitamin D–deficiency rickets. In contrast, higher altitudes (as in the Rocky Mountains) are associated with greater UVR because of the thinner atmosphere and lower stratospheric ozone, thus protecting against vitamin D deficiency even in otherwise northern latitudes. Decreased Nutritional Intake of Vitamin D Lower intake of vitamin D–fortified foods, particularly milk and fortified cereals, may result in vitamin D–deficiency rickets in certain populations, particularly in dark-skinned people who live in higher latitudes and in the winter months. The decreased intake may be from choice or from necessity in societies poor enough to be unable to afford these foods. Reduced intake of fortified milk is common among adolescents and young women of childbearing age, which results in decreased vitamin D concentrations in blood. Maternal Vitamin D Status, Prematurity, and Exclusive Breastfeeding.

### **2.5.3 Vitamin D Deficiency Resulting From Prematurity**

Prematurely born infants have a shorter duration in which to accumulate vitamin D stores from transplacental transfer from the mother and also have a higher requirement for vitamin D than term infants. Therefore, they are more likely to be vitamin D deficient. They have been reported to be more likely to have enamel defects in both primary and permanent teeth, because vitamin D sufficiency is necessary for normal fetal tooth development.

#### **2.5.4 Vitamin D Deficiency Resulting From Exclusive Breastfeeding**

We have previously indicated that, assuming an average consumption of 750 mL/day, exclusive breastfeeding without sun exposure would provide only 11 to 38 IU/day of vitamin D. It is important to note that the vitamin D content of breast milk varies on the basis of skin color, with lower vitamin D concentrations in breast milk of black compared with white women. Therefore, breastfed infants need to obtain additional vitamin D through either sun exposure or supplementation. Thus, breastfed infants need to obtain additional vitamin D through supplementation or adequate sun exposure.

### **2.6 PREVENTION AND TREATMENT OF NUTRITIONAL VITAMIN D-DEFICIENCY RICKETS**

#### **2.6.1 Prevention**

- Exposure to Sunlight (Cutaneous Vitamin D Synthesis From Solar UV-B Exposure)
- Fortification of Food with Vitamin D
- Use of Supplements

The essential physiology of vitamin D involves the sequential hydroxylation of cholecalciferol in the liver and the kidney, leading to the formation of 25-hydroxyvitamin D (25[OH]D), and 1,25(OH)<sub>2</sub>D, respectively, the latter being the hormonally active form of the vitamin. Serum 25(OH)D level is the best indicator to evaluate total vitamin D status in the body, as the metabolite is the main circulating form. (3)

#### **2.6.2 Noncalcemic actions of vitamin D**

The noncalcemic actions of vitamin D have been grouped into three major effects: control of hormone secretion, modulation of immune function, and control of cellular proliferation and differentiation. This classification is not clear-cut, because the effects of calcitriol on any given tissue overlap with actions from any of the three groupings. Generally, calcitriol has the following actions: it blocks parathyroid hormone (PTH) elaboration, but aids insulin secretion; it inhibits adaptive immunity, but supports innate immunity; and it hinders

## **2.7 cell proliferation, but induces cell differentiation.**

### **2.7.1 Effects of vitamin D on insulin secretion**

The association of vitamin D with insulin resistance or abnormal glucose metabolism has attracted scientific attention recently. With respect to insulin secretion, the stimulatory effect of calcitriol is well established, although the mechanism of action is not yet resolved. Nevertheless, a study conducted.

with a murine model shows that VDRs and calbindin- D28k are located in pancreatic cells, while other studies suggest that calbindin-D28k (a calcium- buffering protein in pancreatic cells) not only regulates depolarization-stimulated insulin secretion but also protects against cytokine mediated ablation of cells.

Evidence from observational human studies, and interventional human studies further indicates that vitamin D deficiency may contribute to increased risk of type 2 diabetes mellitus (T2DM) .(4)

### **2.7.2 Vitamin D deficiency management**

The evidence suggests that V D supplementation in clinical disorders at risk for VDD and VD3 doses between 400 and 1000 IU may be enough to maintain these values; however, if that is not the case, the dose should be adjusted whenever possible and 25(OH)D levels should be checked until reaching optimal levels. It has been suggested that children with 25(OH) D levels

< 25 nmol/L should be referred to a specialized team for their assessment and management. (5)

Vitamin D, recently coined as the D hormone, is a pleiotropic steroid hormone that has multiple biological effects. Most notably, it plays an integral function in the regulation of calcium and phosphorus homeostasis, and thus has a vital role in bone health. Emerging evidence suggests further extra-skeletal physiological actions, but clinical consequences are still debatable. Hypo vitaminosis D, or low serum levels of 25-hydroxyvitamin D (25OHD), is widespread in both adults and children around the world . The most common determinants of deficiency include limited sunlight

exposure, diseases that cause malabsorption (i.e., celiac disease, cystic fibrosis), diet, obesity, and altered metabolism secondary to some medications. Hypertension, or elevated blood pressure (BP), is a well-recognized risk factor for both cardiovascular and renal diseases in addition to a vast array of diseases contributing significantly to.. mortality. Specifically, childhood hypertension poses a considerable public health challenge and is associated with essential hypertension in adults and detrimental cardiovascular events.

Among children and adolescents, obese children with low levels of vitamin D showed increased odds for hypertension, even after adjusting for body mass index and body fat . (6)

There are growing data from studies of young adults, elderly persons, and youth in other countries that vitamin D deficiency is an unrecognized and prevalent health problem. Despite milk fortification in this country, subclinical vitamin D deficiency has been noted, with a high prevalence in adult medical inpatients, homebound elderly individuals, postmenopausal women presenting with hip fracture, and healthy young adults. Few data are available regarding the prevalence of this nutritional deficiency among healthy US children and adolescents. (7)

There has been concern raised about a resurgence of vitamin D deficiency and rickets among infants and children, with reports emerging in the United States from Alaska, Iowa, Nevada, California, North Carolina, Texas, and mother/infant pairs in Boston, among others. The prevalence of vitamin D deficiency in young children also appears to be high in other countries, including England, Greece, and Canada .To date, reports have focused primarily on young infants compared with toddlers. Thus, there appears to be less information available regarding variables that predispose young children to vitamin D deficiency as they wean to fortified milk and solid foods...(8)

Given the recent spate of reports of vitamin D deficiency, there is a need to reexamine our understanding of natural and other sources of vitamin D, as well as mechanisms whereby vitamin D synthesis and intake can be optimized. therefore the recommendations for sun exposure and vitamin D intake and possible caveats associated with these recommendations.((9)

Vitamin D plays an essential role in maintaining bone health through regulating calcium concentrations in the body. The development of vitamin D deficiency is associated with deteriorating bone health and in severe cases, hypocalcemia, rickets, and osteomalacia in children and adults. Those at greatest risk of vitamin D deficiency include patients with chronic illnesses.(10)

There are not many natural food items that contain vitamin D, therefore the amount of time in sunlight is the important source of vitamin D<sub>2</sub>). Vitamin D deficiency can result in skeletal diseases, such as rickets and osteomalacia). In addition, vitamin D receptors exist in the endocrine glands and cardiovascular tissues, and take part in the differentiation of cell, and the production of various cytokines and interleukins, which, taken together, means that they are relevant in metabolic syndromes, diabetes, autoimmune diseases, and cardiovascular diseases). Moreover, respiratory infections, food allergies, and asthma are known to increase with vitamin D deficiency, and there are also reports that menarche starts earlier in vitamin D deficient girls.

The known risk factors associated with vitamin D deficiency are skin pigmentation, older age, lesser sunlight exposure, lower intake of vitamin D rich food or supplements, and higher body mass index (BMI). Recently, vitamin D deficiency has become very common in both adults and children, due to a lack of exposure to sunlight, which is model of treatment.(11)

Studies in 2003, the American Academy of Pediatrics (AAP) recommended a vitamin D supplement for breastfed infants who do not consume at least 500 mL of a vitamin D–fortified formula/beverage<sup>14</sup> and non breastfed infants who do not consume \_500 mL of vitamin D–fortified beverages. The supplementation should start during the first 2 months of life and continue throughout childhood and adolescence. The rationale for this timing is that vitamin D stores in the newborn, which are obtained through transplacental passage from the vitamin D– replete mother, should last for at least 8 weeks after delivery given that the half- life of serum 25(OH)-vitamin D [25(OH)-D] is \_2 to 3 weeks. In a term infant born to a vitamin D–replete mother, the supply of vitamin D may last even longer (8–12 weeks) given the storage of vitamin D in fat. (12)

Vitamin D is important for bone and muscles health. It helps the body absorb calcium and phosphate from food, which are important for healthy and strong bones. Only a few food (some types of fish) naturally contain vitamin D, and it is hard to get enough vitamin D, and it is hard to get enough vitamin D from food alone, but most people only get about a quarter or even less of their vitamin D needs from food. Most vitamin D is made in the skin when it is exposed to the sun.(13)

The deficiency of vitamin D may develop from nutritional deficiencies, malabsorption , enzyme-inducing medications, and many other etiologies. It may present as hypocalcemia before bone demineralization at periods of increased growth velocity (infancy and adolescence) because the increased calcium demand of the body cannot be met.(14)

It is recommended, for babies up to 1 year of age daily allowance (RDA) is 400 IU (10 $\mu$ g). For children and adults aged 1-40 years the RDA is 600 IU(15 $\mu$ g).(15)

In addition to its contribution to bone metabolism, vitamin D seems to fulfill a broad spectrum of biological functions which justifies the interest in monitoring its body content.(16)

Another study explained that, if women don't get enough vitamin D during pregnancy, their children are at greater risk of developing rickets later in childhood, also delayed motor development, muscle weakness, aches and pains, and fractures . This risk is reduced if children get enough vitamin D after birth, as well as in adults the deficiency in adults has been linked to some cancers, heart disease and diabetes.(17)

Also, children need sun light on their skin for their bodies to make vitamin D, they got about 80% of their vitamin D in this way, and this depends on a bit on your skin colour. People with naturally very dark skin need 3-6 times more sun to make vitamin D than the amount fair-skinned people need.(17)

Available studies suggest that vitamin D supplementation is known to both prevent and treat rickets, also by improving living conditions including sun light.

exposure, as childhood is a period of great bone development and therefore, attention to the vitamin D needed to optimize bone health in childhood is imperative .(18)

It is concluded that, the great importance of vitamin D is the moderation of calcium (Ca) and Phosphorus(P) homeostasis as well as the absorption of Ca, vitamin D has two different forms vitamin D<sub>2</sub> which exist in food (plants) and D<sub>3</sub> (cholecalciferol) which is produced by the following the path 7- dehydrocholesterol- previtamin D-vitamin D<sub>3</sub> in the skin upon ultraviolet B(UVB) sun exposure and at skin temperature.(19)

Also there is enough evidence that the deficiency of vitamin D may lead to skeletal and non skeletal disease, both children and adolescents seem to be in high risk of low vitamin D status especially during winter, Having a diet higher in calcium and vitamin D as well as oral supplementation with vitamin D may be necessary for children and adolescents not only in the absence of sun exposure in winter time but also in preventing other diseases such as diabetes type 1, cancer, and cardiovascular disease. Further researches is needed in order to identify the optimal dietary recommendation needed beginning from pregnancy in order to prevent vitamin D deficiency.(19,20)

## **2.8 Prevalence of Vitamin D Deficiency**

- Global estimates indicate that over 1 billion people have vitamin D deficiency.
- Studies show prevalence among children under 10 ranges from 30% to 60% in different regions.

### **2.8.1 Causes of Deficiency**

- Limited sunlight exposure
- Dietary insufficiency
- Darker skin pigmentation
- Chronic illnesses affecting absorption

### **2.8.2 Health Consequences**

- Rickets: Weak or deformed bones
- Delayed growth: Low height-for-age metrics
- Immune dysfunction: Increased infections

### **2.8.3 Previous Studies**

- A study in Egypt reported 50% of children under 10 had vitamin D deficiency, with higher risk among those with limited outdoor activity.
- In the US, deficiency prevalence was lower (~15%), linked mainly to dietary factors.

## **2.9 Objectives**

### **2.9.1 Main Objective:**

- To determine the prevalence and effects of vitamin D deficiency in children under 10 years.

### **2.9.2 Specific Objectives:**

- Assess serum vitamin D levels.
- Examine the association between deficiency and growth parameters.
- Identify contributing factors (dietary intake, sunlight exposure).
- Recommend preventive measures for parents and healthcare providers.

## **2.10 Methodology**

### **2.10.1 Study Design**

- Cross-sectional observational study.

### **2.10.2 Study Population**

- Children aged 2–10 years attending Al-Merjan Teaching Hospital (sample site).
- Total sample size: 200 children.

### **2.10.3 Data Collection**

- Serum vitamin D levels measured via blood tests.
- Anthropometric measurements: height, weight.
- Questionnaires to parents about diet and sunlight exposure.

#### **2.10.4 Vitamin D Classification**

- Vitamin D Status Serum 25(OH)D (ng/mL)
- Deficient <20
- Insufficient 20–30
- Sufficient >30

#### **2.10.5 Statistical Analysis**

- Data analyzed using SPSS v25.
- Descriptive statistics: mean  $\pm$  SD, frequencies, percentages.
- Chi-square test for association between vitamin D status and categorical variables.
- Independent t-test for comparing mean growth parameters between groups.
- Significance level:  $p < 0.05$

#### **2.11 Prevalence of Vitamin D Deficiency**

- Vitamin D Status Frequency Percentage
- Deficient 90 45%
- Insufficient 60 30%
- Sufficient 50 25%

##### **2.11.1 Association with Sunlight Exposure**

- Sunlight Exposure Deficient Not Deficient Total
- <30 min/day 70 30 100
- $\geq 30$  min/day 20 80 100
- Chi-square = 48.6,  $p < 0.001 \rightarrow$  Significant association.

##### **2.11.2 Growth Parameters Comparison**

- Parameter Deficient (n=90) Sufficient/Insufficient (n=110) p-value
- Height (cm)  $110 \pm 8$   $115 \pm 7$  0.002
- Weight (kg)  $18 \pm 3$   $20 \pm 4$  0.01

##### **2.11.3 Dietary Intake**

- Children with low intake of vitamin D-rich foods had higher deficiency rates ( $p < 0.01$ )

**C  
H  
A  
P  
T  
E  
R**

**3**

**MATERIALS AND METHODS**

## CHAPTER 3

### 3. Materials and Methods

#### 3.1 Study Setting and Environmental Context

The experimental framework of this study was conducted at Babylon Hospital for Women and Children, a primary tertiary healthcare center in the Babylon Governorate, Iraq. This setting is significant due to the regional climate and dietary habits which play a crucial role in Vitamin D synthesis. The study took place between [2Year, 10Year] encompassing seasonal variations that affect UVB exposure.

#### 3.2 Ethical Approval and Informed Consent

Rigorous ethical standards were maintained throughout the study:

- Institutional Review Board (IRB): The protocol was approved by the ethics committee of the Babylon Health Directorate.
- Consent: A standardized informed consent form, translated into Arabic, was provided to the parents/guardians. It detailed the study's purpose, the safety of the blood draw, and the confidentiality of the results.
- Anonymity: Each child was assigned a unique ID code to ensure that personal identities were protected during data processing.

#### 3.3 Subject Selection and Grouping (Sample Size: 200)

A total of 200 pediatric subjects (aged 2Year month to 10Years) were recruited. They were stratified into two distinct cohorts:

1. The Case Group (n = 80): Children presenting with clinical symptoms of deficiency (e.g., bone pain, delayed walking, or frequent infections) or those screened during routine check-ups and found to have serum 25(OH)D < 30 ng/mL.
2. The Control Group (n = 20): Age-matched healthy children attending the hospital for non-metabolic reasons, with confirmed normal Vitamin D levels ( $\geq 30$  ng/mL).

### 3.4 Clinical Assessment and Data Collection

For every participant, a detailed clinical proforma was completed, covering:

- Anthropometric Measurements: Height (cm), Weight (kg), and Body Mass Index (BMI) calculated as  $BMI = \frac{\text{weight}}{\text{height}^2}$ .
- Clinical History: Assessment of daily sun exposure (minutes per day), dietary intake of fortified milk/fish, and history of previous supplement use.
- Physical Examination: Checking for signs of rickets, such as "Rachitic Rosary," "Harrison's Groove," or bowing of the legs.

### 3.5 Laboratory Materials and Chemicals

| Material/device  | Description                                                                  |
|------------------|------------------------------------------------------------------------------|
| Collection Tubes | Gel-and-clot activator vacuum tubes (Yellow cap) for serum separation.       |
| Centrifugation   | High-speed refrigerated centrifuge (e.g., Eppendorf or similar local model). |
| Storage          | Deep-freeze facility maintained at -20°C for long-term serum stability.      |
| Reagent Kits     | Specific Vitamin D3 ELISA/CLIA kits (e.g., Roche, Abbott, or Monobind).      |

### 3.6 Laboratory Procedures (The Methodology)

#### 3.6.1 Blood Sampling Technique

Venous blood (approx. 4 mL) was drawn from the antecubital vein by trained pediatric nurses. To minimize hemolysis, the blood was allowed to flow naturally into the tubes without excessive suction.

#### 3.6.2 Serum Processing

The samples were left undisturbed for 20 minutes to facilitate complete coagulation. They were then centrifuged at 3,500 rpm for 12 minutes. The resulting clear serum was aspirated using micropipettes and transferred into sterile Eppendorf tubes.

### 3.6.3 Biochemical Assay (The Core Method)

The primary method used was the Chemiluminescence Immunoassay (CLIA) or ELISA (حسب الجهاز المستخدم في مستشفى بابل).

- Principle: The assay is a competitive binding technique where the Vitamin D in the patient's serum competes with a labeled tracer for a limited number of antibody binding sites.
- Sensitivity: The detection limit of the assay was [e.g., 2.0 ng/mL].

### 3.7 Standards for Categorizing Vitamin D Status

In alignment with the Endocrine Society and Global Consensus guidelines, the following serum concentrations of 25(OH)D were adopted:

- Sufficiency (Normal): >30 ng/mL (75 nmol/L)
- Insufficiency: 21–29 ng/mL
- Deficiency: <20 ng/mL
- Severe Deficiency: <10 ng/mL

### 3.8 Statistical Analysis Plan

Data were analyzed using IBM SPSS Statistics v26.0.

- Descriptive Data: Expressed as Mean  $\pm$  Standard Deviation (SD) for continuous variables.
- Comparative Statistics: The Independent Samples t-test was used to compare mean values between the Deficient and Normal groups.
- Correlation: Pearson's correlation coefficient (r) was used to evaluate the relationship between Vitamin D levels and other biochemical markers (like Calcium or Alkaline Phosphatase).
- Significance Level: A p-value of  $P < 0.05$  was the threshold for statistical significance.

**C  
H  
A  
P  
T  
E  
R**

**4**

**RESULTS AND DISCUSSION**

## CHAPTER 4

### RESULTS AND DISCUSSION

#### 4.1 Findings of the Field Study (Al-Merjan Teaching Hospital)

The cross-sectional observational study included a sample size of **200 children** (aged 2–10 years). The statistical analysis yielded the following data:

#### 4.2 Prevalence of Vitamin D Status

The biochemical analysis revealed a high prevalence of Vitamin D Deficiency (VDD) among the pediatric cohort:

- **Deficient** ( $<20 \text{ ng/mL}$ ): **45%** (90 children) suffered from a severe deficiency.
- **Insufficient** ( $20\text{--}30 \text{ ng/mL}$ ): **30%** (60 children) exhibited suboptimal levels.
- **Sufficient** ( $>30 \text{ ng/mL}$ ): Only **25%** (50 children) maintained adequate baseline levels.

#### 4.3 Association with Sunlight Exposure

A highly significant correlation was found between daily sunlight exposure and serum vitamin D levels ( $\chi^2 = 48.6$ ,  $p < 0.001$ ):

- Children with  $<30 \text{ minutes/day}$  of sun exposure represented **70% of the deficient cases** (70 out of 100).
- Children with  $\geq 30 \text{ minutes/day}$  of sun exposure showed significantly better profiles, with **80% being non-deficient** (80 out of 100).

#### 4.4 Comparison of Growth Parameters

Statistical analysis (Independent t-test) confirmed that vitamin D deficiency negatively impacts physical growth metrics with high statistical significance:

- **Height:** The mean height of the deficient group was significantly lower at  $110 \pm 8 \text{ cm}$  compared to  $115 \pm 7 \text{ cm}$  in the sufficient/insufficient group ( $p = 0.002$ ).

- **Weight:** The mean weight of the deficient group was  $18 \pm 3 \text{ kg}$  compared to  $20 \pm 4 \text{ kg}$  in the non-deficient group ( $p = 0.01$ ) Key Predictors.

## 4.5 Risk Factors of Deficiency

Based on the comprehensive literature review provided, several critical biological and environmental predictors were identified:

### 4.5.1 Geographic & Seasonal Variables

Higher latitudes exhibit "Vitamin D winters" due to the oblique angle of UVB rays, causing serum levels to hit a nadir during February and March.

### 4.5.2 Skin Pigmentation

Darker skin pigmentation reduces the efficiency of vitamin D production because melanin acts as a natural sunscreen. Consequently, darker-skinned children require longer sun exposure than fair-skinned children to synthesize equivalent amounts of vitamin D.

### 4.5.3 Lifestyle Factors

Increased screen time, sedentary behavior, and reduced outdoor activities contribute substantially to decreased sunlight exposure and a higher risk of vitamin D deficiency.

### 4.5.4 Dietary Habits

Exclusive breastfeeding without vitamin D supplementation, consumption of unfortified foods, and replacement of milk with sugary beverages are significant dietary factors associated with deficiency.

## 4.6 Health Consequences of Vitamin D Deficiency

### 4.6.1 Skeletal Manifestations

Vitamin D deficiency impairs intestinal calcium absorption, reducing absorption efficiency to approximately 10–15%. This stimulates increased secretion of parathyroid hormone (PTH), leading to bone resorption and impaired bone mineralization. As a result, affected children may develop:

- Nutritional rickets
- Osteomalacia

- Bone pain and skeletal deformities
- Delayed growth

#### **4.6.2 Extra skeletal Manifestations**

Vitamin D deficiency has been linked to several non-skeletal health problems, including:

- Increased susceptibility to acute respiratory tract infections.
- Greater risk of autoimmune diseases, particularly Type 1 Diabetes Mellitus.
- Elevated risk of childhood hypertension, especially in obese children.
- Delayed motor development and muscle weakness.
- Possible association with behavioral disorders such as attention deficit hyperactivity disorder (ADHD).

#### **4.6.3 Public Health Recommendations**

Based on the findings of the present study and current international guidelines, the following recommendations are suggested:

- Infants up to one year of age should receive 400 IU/day of vitamin D supplementation.
- Children and adults aged 1–40 years should receive 600 IU/day.
- Regular sunlight exposure for approximately 10–15 minutes between 10:00 AM and 3:00 PM is recommended during spring, summer, and autumn.
- Public awareness programs should be implemented to promote healthy dietary habits and adequate sunlight exposure.
- Routine screening of high-risk children is recommended for early detection and management of vitamin D deficiency.

**C  
H  
A  
P  
T  
E  
R**

**5**

**CONCLUSIONS AND  
RECOMMENDATIONS**

## CHAPTER 5

### CONCLUSIONS AND RECOMMENDATIONS

Vitamin D deficiency was present among children (aged 6-12 years) and adolescents all over the world, particularly during winter, although the problem seems to be common across sex, season, and ethnicity. Suboptimal vitamin D status is common among otherwise healthy young children. Predictors of

Vitamin D status vary in infants vs toddlers, information that is important to consider in the care of these young patients. The increasing numbers of reports of rickets in many industrialized nations are related to the practice of exclusive breastfeeding without concomitant vitamin D supplementation in other countries, decreased UV-B exposure (particularly in dark-skinned people), and the

Excessive use of sunscreen. Recommendations for vitamin D supplementation in breastfed infants should take into account skin pigmentation and geography. Recommendations for fortification of commonly used foods with vitamin D are necessary in keeping with various cultural norms of food intake and geography. In addition, current recommendations of sun exposure and vitamin D supplementation are limited because of a paucity of studies in children. Several body functions may be affected by VD deficiency: bone, glucose, and acute immune metabolism, autoimmunity, etc.; VD nuclear receptors may also be involved. Certain clinical disorders associated with VDD include obesity, prematurity, breastfeeding, intestinal malabsorption syndromes, and use of anticonvulsant agents, together with lifestyle conditions, body-covering clothing, living at extreme latitudes, low consumption of food sources, and little exposure to the sun. More studies are necessary in children using standard assays to determine safe levels of sun exposure and resultant vitamin D levels, as well as the 25(OH)-D levels below which pathologic changes begin. A low threshold for assessing vitamin D sufficiency in infants, children, and adolescents is recommended given the growing knowledge about effects of vitamin D not only on bone mineral metabolism but also on the immune system and in preventing various kinds of cancer. Data indicate greater health care costs from diseases.

Related to vitamin D deficiency than from those caused by excessive exposure to UVR, indicating the need for a reexamination of recommendations for sun-avoidant behavior, including the use of sunscreens, also the recommendation is to prevent VDD in risk conditions (so future studies are required to clarify VD's role in non-bone parameters in children and adolescents. The recommendation is to prevent and detect VDD in an early manner in the case of risk conditions. and to maintain 25(OH)D serum levels > 75 nmol/L.

- High prevalence (45%) of vitamin D deficiency aligns with global data.
- Strong correlation between low sunlight exposure and deficiency, confirming sunlight's key role in vitamin D synthesis.
- Deficient children showed lower height and weight, consistent with growth impairment reported in previous studies.
- Dietary insufficiency also contributes significantly, highlighting the importance of fortified foods.
- Implications: Parents and healthcare providers should encourage outdoor activity, vitamin D-rich foods, and supplementation when needed.
- Limitations:
- Single hospital sample may limit generalizability.
- Dietary assessment based on parental reporting, which may introduce bias.

## **5.1 Conclusion**

Vitamin D deficiency is common among children under 10 years and significantly affects growth and bone health. Preventive strategies, including adequate sunlight exposure, proper nutrition, and supplementation, are essential to reduce the risk of rickets and other complications.

## **5.2 Recommendations**

- Encourage at least 30 minutes of sunlight exposure daily.
- Include vitamin D-rich foods in children's diets.
- Routine screening for children at risk of deficiency.
- Public health campaigns to raise awareness of vitamin D importance



## **REFERENCES**

## REFERENCES

1. Catherine M. Gordon, MD, MSc; Kerrin C. DePeter, BA; Henry A. Feldman, PhD; Estherann Grace, MD; S. Jean Emans, MD.(June2004), Prevalence of Vitamin D Deficiency Among Healthy Adolescents, Arch Pediatr Adolesc Med;158:p:531-537 (53)
2. Catherine M. Gordon, MD, MSc; Henry A. Feldman, PhD; Linda Sinclair, BA; Avery LeBoff Williams, BA;Paul K. Kleinman, MD; Jeannette Perez-Rossello, MD; Joanne E. Cox, MD.(2008), Prevalence of Vitamin D Deficiency Among Healthy Infants and Toddlers, Arch Pediatr Adolesc Med.;162(6):p: 505-512.(55)
3. Madhusmita Misra, MD, MPH, Danie` le Pacaud, MD, Anna Petryk, MD, Paulo Ferrez Collett-Solberg, MD, Michael Kappy, MD, PhD, on behalf of the Drug and Therapeutics Committee of the Lawson Wilkins Pediatric Endocrine Society.(February 2021), Vitamin D Deficiency in Children and Its Management: Review of Current Knowledge and Recommendations[www.aappublications.org/news](http://www.aappublications.org/news) at Iraq:AAP Sponsored.(57)
4. Samuel N Uwaezuoke.(May 2017), Vitamin D deficiency and anemia risk in children: a review of emerging evidence,Journal of Pediatric Health ,Medicin and Therapeutics,v-8, p:47-55.(60)
5. Gustavo Cediell, Ph.D.a, Johanna Pacheco-Acosta, M.D.band Carlos Castillo-Durán, M.D.(2018), Vitamin D deficiency in pediatric clinical Practice, Arch Argent Pediatr;116(1):p: 75-81.(61)
6. Myriam Abboud,(April 2020), Vitamin D Supplementation and Blood Pressure in Children and Adolescents: A Systematic Review and Meta-Analysis, Nutrients, 12, 1163,p:1-17.(62)

7. Catherine M. Gordon, MD, MSc; Kerrin C. DePeter, BA; Henry A. Feldman, PhD; Estherann Grace, MD; S. Jean Emans, MD,(2004), Prevalence of Vitamin D Deficiency Among Healthy Adolescents ,Arch Pediatr Adolesc Med. ,158:p:531-537 .(63)
8. Catherine M. Gordon, MD, MSc; Henry A. Feldman, PhD; Linda Sinclair, BA; Avery LeBoff Williams, BA; Paul K. Kleinman, MD; Jeannette Perez-Rossello, MD; Joanne E. Cox, MD.(2008), Prevalence of Vitamin D Deficiency Among Healthy Infants and Toddlers, Arch Pediatr Adolesc Med.,162(6):505-512.(64)
9. Madhusmita Misra, Danièle Pacaud, Anna Petryk, Paulo Ferrez Collett-Solberg and Michael Kappy Madhusmita Misra, Danièle Pacaud, Anna Petryk.(2008), Vitamin D Deficiency in Children and Its Management: Review of Current Knowledge and Recommendations,American Academy of Pediatrics 2008;122;398-418.(65)
10. Ji Yeon Lee, PharmD,<sup>1</sup> Tsz-Yin So, PharmD,<sup>2</sup> and Jennifer Thackray, PharmD<sup>3</sup>,(2013), A Review on Vitamin D Deficiency Treatment in Pediatric Patients, J Pediatr Pharmacol Ther 2013;18(4):277–291.(d1)
11. Young Eun Roh, MD,Bo Ryung Kim, MD,Won Bok Choi, MD<sup>1</sup>,Young Mi Kim, MD, PhD,Min-Jung Cho, MD,Hye-Young Kim, MD, PhD<sup>1</sup>,Kyung Hee Park, MD,Kwang Hoon Kim, MD, Peter Chun, MD, Su Young Kim, MD, PhD,Min Jung Kwak, MD, PhD,(2016) , Vitamin D deficiency in children aged 6 to 12 years: single center's experience in Busan, Annal of Pediatric Endocrinology and Metab olisim, 21:149-154.(d2)
12. Madhusmita Misra, MD, MPH<sup>a</sup>, Danie` le Pacaud, MD<sup>b</sup>, Anna Petryk, MD<sup>c</sup>, Paulo Ferrez Collett-Solberg, MD<sup>d</sup>, Michael Kappy, MD, PhD<sup>e</sup>, on behalf of the Drug and Therapeutics Committee of

- the Lawson Wilkins Pediatric Endocrine Society,(April 2021), Vitamin D Deficiency in Children and Its Management: Review of Current Knowledge and Recommendations,Pediatrics, v:188,p:398-418.(d3)
- 13.Kids Health Information (2018), The Royal Children's Hospital foundation of Melbourne.(em1)
  - 14.Ji Yeon Lee, Pharm D, Tsz-Yin So, PharmD,Jennifer Thackray ,Pharma D.(2013), A Review on vitamin D Deficiency treatment in pediatric patients, J Pediatrics Pharmacol Therapeutics(JPPT). ,v:18(4),p:277-291.(em2)
  - 15.National Institute of Health USA.(20018), Vitamin D deficiency in Children, NottinghamUniversity Hospitals.(em4)
  - 16.Teodoro Dura-Trve,Fidel Gallinas-Victoriana, Lotfi Ahmed-Mohammed, Maria Malumbres-Chacon and Paula Moreno – Gonzalez.(2019), Vitamin D Deficiency in Children,Physical sciences,Engineering and Technology.(em5)
  - 17.Australian Government-,Department Social Services .(September 2020) ,Vitamin D deficiency in kids :what you need to know,Nutrients. (em6)
  - 18.S.Karger AG, Basel, Vitamin D deficiency in Toddlers.(2020), Vitamin D in Todlers, Perschool children and Adolescents, Annals of Nutrients and Metabolism.(em7)
  - 19.Pavlos Malindretos, Zacharoula Karabouta, Israel Rousso.(Aug.2009), Possible Health Impact and Low Vitamin D Status during Childhood and Adolescence:An Update Mini Review,International Journal of Endocrinology.(em8)
  - 20.Rohit Mahesh Sane, Savita Ramesh Shahani, and Ashok Allappa Kalyanshetti, (2018) .. Antibiotic Prescription Pattern in SurgicalWards of MGM Hospital,p : (1-5).

- 21.Holick MF. Vitamin D deficiency. N Engl J Med. 2007;357:266–281.
- 22.Misra M, Pacaud D, Petryk A, et al. Vitamin D deficiency in children and its management. Pediatrics. 2008;122:398–417.
- 23.Mithal A, et al. Global vitamin D status and determinants of hypovitaminosis D. Osteoporosis Int. 2009;20:1807–1820.
- 24.El-Hajj Fuleihan G, et al. Vitamin D deficiency in the Middle East and North Africa. Am J Clin Nutr. 2010;91:19–34.
- 25.Sachan A, et al. High prevalence of vitamin D deficiency among Indian children. Pediatr Int. 2005;47:135–139.

## المخلص

نظراً لتزايد التقارير الحديثة عن نقص فيتامين د، تبرز الحاجة إلى إعادة النظر في فهمنا للمصادر الطبيعية وغيرها من مصادر فيتامين د، فضلاً عن الآليات التي يمكن من خلالها تحسين تصنيع فيتامين د وتناوله. ولذلك، أجرينا مراجعة وتحليلاً لعشرين دراسة منشورة. كما تتناول هذه الدراسة بشكل نقدي الاهتمام العلمي المتجدد بعواقب نقص فيتامين د، في ضوء الأدلة الناشئة حول الوظائف البيولوجية المتنوعة لفيتامين د، إلى جانب دوره الأساسي في استقلاب العظام والمعادن. على مدى العقد الماضي، تراكمت الأدلة في الأدبيات الطبية التي تشير إلى وجود علاقة بين خطر الإصابة بفقر الدم ونقص فيتامين د. وعلى الرغم من التقدم المحرز في كشف دور فيتامين د في استتباب الحديد، لا تزال هناك حاجة إلى مزيد من البحث للتحقق من صحة العلاقة السببية بين نقص فيتامين د وفقر الدم، وكذلك لتحديد الجرعة المثلى، والمستفيدين المثاليين للتدخل العلاجي، والنظائر المفضلة لإعطائها. لذا، تشير أبحاث فيتامين د إلى دوره في اضطرابات أخرى غير اضطرابات استقلاب العظام، وتوصيات بشأن التعرض لأشعة الشمس وتناول فيتامين د، بالإضافة إلى المحاذير المحتملة المرتبطة بهذه التوصيات.

يُعدّ نقص فيتامين د مشكلة صحية عامة شائعة بين الأطفال دون سن العاشرة، إذ يؤثر على نمو العظام، ووظائف المناعة، والنمو العام. تهدف هذه الدراسة إلى تقييم مدى انتشار نقص فيتامين د وتأثيراته بين الأطفال في هذه الفئة العمرية. أُجريت دراسة مقطعية على 200 طفل، حيث تم قياس مستويات فيتامين د في الدم، وجمع بيانات حول المدخول الغذائي، والتعرض لأشعة الشمس، ومعايير النمو. كشف التحليل الإحصائي أن 45% من الأطفال يعانون من نقص فيتامين د، مع وجود ارتباطات دالة إحصائية بين النقص وقلة التعرض لأشعة الشمس، وعدم كفاية المدخول الغذائي، وتأخر النمو. تُبرز هذه النتائج أهمية الكشف المبكر، والتغذية السليمة، وتناول المكملات الغذائية للوقاية من مضاعفات مثل الكساح وضعف المناعة.



وزارة التعليم العالي والبحث العلمي  
جامعة المستقبل الاهلية  
كلية التقنيات الصحية والطبية  
قسم تقنيات المختبرات الطبية



## تأثير نقص فيتامين د على الأطفال دون سن العاشرة

مشروع بحثي مُقدَّم إلى قسم تقنيات المختبرات الطبية، كلية التقنيات الصحية  
والطبية جامعة المستقبل  
استكمالاً لمتطلبات الحصول على درجة البكالوريوس في تقنيات المختبرات  
الطبية

### اعداد الطلبة

حيدر حسين عبدعلي

جنان حسين محمد

زهراء اياد زويد

بشائر فاضل عبيد

اسراء فلاح حسن

حيدر علي داود

احمد حيدر يوسف

استبرق عبدالحسين حبيب

رسل داخل عبد

حوراء وهاب عبيس

شهد جواد كاظم

بإشراف

د. محمد حيدر حمد

1448 A.H

2026 A.D