

ALMUSTAQBAL UNIVERSITY COLLEGE

Biomedical Engineering Department

Stage : Second year students

Subject : Chemistry 1 - Lecture 6

Lecturer: Assistant professor Dr. SADIQ . J. BAQIR



Molarity of liquids:

The molarity of liquids Can be determined by applying the following formula:

$$\text{Molarity of liquid (M)} = \frac{\text{sp.gr} \times \left(\frac{w}{w}\right)\% \times 1000}{Mwt}$$

$$\text{Specific gravity (Sp.gr)} = \frac{\text{density of substance}}{\text{density of water}}$$

$$\text{Specific gravity (Sp.gr)} = \frac{d_{\text{substance}}}{d_{\text{H}_2\text{O}}}$$

$$(\text{sp.gr} \approx d_{\text{substance}}) \text{ as } d_{\text{H}_2\text{O}} = 1$$

Example:

Calculate the molarity of the solution of 70.5 % HNO₃ (w/w) (63.0 g/mol) that has specific gravity of (1.42) .

Solution:

$$\text{Molarity (M)} = \frac{\text{sp.gr} \times \left(\frac{w}{w}\right)\% \times 1000}{Mwt}$$

$$M = \frac{1.42 \times \left(\frac{70.5}{100}\right) \times 1000}{63.0} = \frac{1.42 \times 70.5 \times 10}{63.0} = 15.9 \text{ M}$$

Example :

Calculate the molarity of NaOH (40 g/mol) solution of 50 $\left(\frac{w}{w}\right)\%$ knowing that its specific gravity(sp.gr) is 1.525 .

Solution:

$$\text{Molarity(M)} = \frac{\text{sp.gr} \times \left(\frac{w}{w}\right)\% \times 1000}{Mwt}$$

$$\text{Molarity (M)} = \frac{1.525 \times \left(\frac{50}{100}\right) \times 1000}{40} = \frac{1.525 \times 50.5 \times 10}{40} = 19.06 \text{ M}$$

Example:

Describe the preparation of (100 mL) of (6.0 M) HCl from its concentrated solution that is 37.1 % (w/w) HCl (36.5 g /mole) and has specific gravity (sp.gr) of (1.181) .

Solution:

١. نحسب تركيز الحامض الاصيلي (المركز) من القانون التالي:

$$M_{\text{HCl}} = \frac{\text{sp.gr} \times \left(\frac{w}{w}\right)\% \times 1000}{Mwt}$$

$$M_{\text{HCl}} = \frac{1.18 \times \frac{37.1}{100} \times 1000}{36.5}$$

$$M_{\text{HCl}} = \frac{1.18 \times 37.1 \times 1000}{36.5 \times 100}$$

$$M_{\text{HCl}} = \frac{1.18 \times 37.1 \times 10}{36.5} = 12.0 \text{ M}$$

The Molarity of the concentrated acid is 12.0M

الآن نذهب الى قانون التخفيف لحساب الحجم المطلوب اخذه من الحامض المركز وتخفيفه الى الحجم المطلوب (١٠٠ مللتر في هذا المثال) وكمايلي:

No. of moles of Conc. solution = No. of moles of dil. Solution

also

No. of mmoles of Conc. solution = No. of mmoles of dil. Solution

$$M_{\text{conc.}} V_{\text{conc.}} = M_{\text{dil.}} V_{\text{dil.}}$$

$$12.0 \times V_{\text{conc}} = 6.0 \times 100$$

$$V_{\text{conc}} = \frac{6.0 \times 100}{12} = 50 \text{ mL.}$$

Then 50 mL of concentrated acid is to be diluted to 100 mL to give 6 M solution

Example:

Describe the preparation of 500 mL of 3.00 M H₂SO₄ (98 g/mol) from the commercial reagent that is 93% H₂SO₄ (w/w) and has a specific gravity of 1.830.

Solution:

1. We have to calculate the concentration of the original conc. Solution

$$M_{\text{H}_2\text{SO}_4} = \frac{\text{sp.gr} \times \% \times 1000}{M.wt}$$

$$M_{\text{H}_2\text{SO}_4} = \frac{1.830 \times \frac{93}{100} \times 1000}{98}$$

$$M_{\text{H}_2\text{SO}_4} = \frac{1.830 \times 93 \times 1000}{98 \times 100}$$

$$M_{\text{H}_2\text{SO}_4} = \frac{1.830 \times 93 \times 10}{98} = 17.37 \text{ M}$$

لحساب الحجم المطلوب اخذه من الحامض المركز وتخفيفه الى الحجم المطلوب (٥٠٠ ملتر في هذا المثال) نطبق قانون التخفيف التالي:

$$M_{\text{conc.}} \cdot V_{\text{conc.}} = M_{\text{dil.}} \cdot V_{\text{dil.}}$$

$$17.37 \times V_{\text{conc}} = 3.0 \times 500$$

$$V_{\text{conc}} = \frac{3.0 \times 500}{17.37} = 86.36 \text{ mL.}$$

Then 86.36 mL of concentrated acid is to be diluted to 500 mL to give 3 M solution.

Calculation of Normality of liquids

$$\text{Normality of liquid (N)} = \frac{\text{sp.gr} \times \left(\frac{w}{w}\right)\% \times 1000}{\text{eq.wt}}$$

Example:

Describe the preparation of 500 mL of 3.00 N H₂SO₄(98 g /mol) from the commercial reagent that is 96% H₂SO₄ (w/w) and has a specific gravity of 1.840.

Solution:

$$M_{\text{H}_2\text{SO}_4} = \frac{\text{sp.gr} \times \% \times 1000}{\text{eq.wt}}$$

$$\text{eq.wt} = \frac{M_{\text{wt}}}{\eta}$$

For H₂SO₄ $\eta=2$ then

$$\text{eq.wt} = \frac{98}{2} = 49$$

$$\text{Normality (N}_{\text{H}_2\text{SO}_4}) = \frac{1.840 \times \frac{96}{100} \times 1000}{49}$$

$$\text{Normality (N}_{\text{H}_2\text{SO}_4}) = \frac{1.840 \times 96 \times 1000}{49 \times 100}$$

$$\text{Normality (N}_{\text{H}_2\text{SO}_4}) = \frac{1.840 \times 96 \times 10}{49} = 36.04 \text{ N}$$

The Normality of the concentrated acid is 36.04 N

لحساب الحجم المطلوب اخذه من الحامض المركز وتخفيفه الى الحجم المطلوب (٥٠٠ ملتر في هذا المثال) نطبق قانون التخفيف التالي:

$$N_{\text{conc.}} V_{\text{conc.}} = N_{\text{dil.}} V_{\text{dil.}}$$

$$36.04 \times V_{\text{conc}} = 3.0 \times 500$$

$$V_{\text{conc}} = \frac{3.0 \times 500}{36.04} = 41.62 \text{ mL.}$$

Then 41.62 mL of concentrated acid is to be diluted to 500 mL to give 3 N solution.

Example:

A solution of 6.42 (w/w)% of $\text{Fe}(\text{NO}_3)_3$ (241.86 g/mol) has a specific gravity of 1.059. Calculate:

- (a) the molar concentration of this solution.
- (b) the mass in grams of $\text{Fe}(\text{NO}_3)_3$ contained in each liter of this solution

Answer:

a) To calculate the molar concentration of the solution

$$M_{\text{Fe}(\text{NO}_3)_3} = \frac{\text{sp.gr} \times \% \times 1000}{\text{Mwt}}$$

$$M_{\text{Fe}(\text{NO}_3)_3} = \frac{1.059 \times \frac{6.42}{100} \times 1000}{241.86} = 0.281$$

(b) the mass in grams of $\text{Fe}(\text{NO}_3)_3$ contained in each liter of this solution (i.e: the concentration of solution in g / L).

Weight (g) = Molarity x volume(liter) x M.wt

$$\text{Weight} = 0.281 \times 1 \text{ liter} \times 241.86 = 67.96 \text{ g}$$

The concentration of solution in g / L = 67.96 g / L

Second method:

$$\text{Molarity}(M) = \frac{\text{wt}(\text{g}) \times 1000}{\text{M.wt} \times V_{\text{mL}}}$$

$$\text{wt}(\text{g}) = \frac{\text{Molarity}(M) \times \text{M.wt} \times V_{\text{mL}}}{1000}$$

$$\text{wt}(\text{g}) = \frac{0.281 \times 241.86 \times 1000_{\text{mL}}}{1000} = 67.96 \text{ g}$$

Example:

A 12.5% (w/w) aqueous solution of NiCl₂ (129.61 g/mol) has specific gravity of 1.149. Calculate:

(a) the Molarity of NiCl₂ in this solution.

(b) the molar concentration of Cl⁻ in the solution.

(c) the mass in grams of NiCl₂ contained in 500 mL of this solution.


Answer:

(a) the Molarity of NiCl₂ in this solution

$$M_{\text{NiCl}_2} = \frac{\text{sp.gr} \times \% \times 1000}{Mwt}$$

$$M_{\text{NiCl}_2} = \frac{1.149 \times \frac{6.42}{100} \times 1000}{129.61} = 0.569 \text{ M}$$

(b) the molarity of Cl⁻ concentration in the solution.



Each 1 mole gives 1 mole 2 mole

Molarity of Cl⁻ = 2 x Molarity of NiCl₂

Molarity of Cl⁻ = 2 x 0.569 = 1.138 M

(c) the mass in grams of NiCl₂ contained in 500 mL of this solution.

Weight (g) = Molarity x volume(liter) x M.wt

$$\text{Weight} = 0.569 \times \left(\frac{500}{1000}\right) \text{ L} \times 129.61 = 36.87 \text{ g}$$

Example:

A solution was prepared by dissolving **327.8 mg** of Na_3PO_4 (163.9 g/mol) in sufficient water to give 750 mL . Calculate:

A) The Molarity and Normality of the solution

B) the Molar concentration of Na^+ in the solution.

Answer:

A) The Molarity and Normality of the solution

$$\text{Molarity(M)} = \frac{\text{wt(g)} \times 1000}{\text{M.wt} \times V_{\text{mL}}}$$

$$\text{Weight of } \text{Na}_3\text{PO}_4 \text{ (g)} = \frac{327.8 \text{ mg}}{1000} = 0.3278 \text{ g}$$

$$\text{Molarity(M)} = \frac{0.3278 \times 1000}{163.9 \times 750} = 0.00267 \text{ M} = 2.67 \times 10^{-3} \text{ M}$$

$$\text{Normality (N)} = \text{Molarity(M)} \times \eta$$

$$(\eta) = \Sigma [\text{no. of cations} \times \text{its valency(cation charge)}]$$

$$\text{For } \text{Na}_3\text{PO}_4 \text{ } (\eta) = \Sigma [3 \text{ Na}^+ \times (+1)] = 3$$

$$\text{Normality (N)} = 2.67 \times 10^{-3} \times 3 = 8.01 \times 10^{-3} \text{ N}$$

B) the Molar concentration of Na^+ in the solution.



1 mole 3 mole

Molarity of Na^+ = 3 x Molarity of Na_3PO_4

$$\text{Molarity of } \text{Na}^+ = 3 \times 2.67 \times 10^{-3} = 8.01 \times 10^{-3} \text{ M}$$