



Al-Mustaqbal University College

Department of Medical Device Technologies

Medical Chemistry

First Stage

Lecture One

Subject:

- Introduction to Analytical Chemistry
- **-Qualitative analysis Chemistry**
- -Quantitative analysis Chemistry

Analytical Chemistry

Introduction

Chemistry: The science deals with the study of matter, including its composition, structure, physical properties, and reactivity. It is divided into five fields:

- 1-Organic chemistry.
- 2-Inorganic chemistry.
- 3-Physical chemistry.
- 4-Biochemical chemistry.
- 5-Analytical chemistry.

Analytical chemistry


Is a measurement science consisting of a set of powerful ideas and methods that are useful in all fields of science, engineering, and medicine.

- * Analytical chemistry concerned with determines the composition of substances. It comprises of two branches:
 - **1-**Qualitative analysis reveals the identity of the elements and compounds in a sample
 - **2-**Quantitative analysis indicates the amount of each substance in a sample.

Chemical Analytical Methods

- Gravimetric methods: Determine the mass of the analyte or some compound chemically related to it.
- Volumetric methods: Measure the volume of a solution containing sufficient reagent to react completely with the analyte.
- Electroanalytical methods: Measure electrical properties such as potential, current, resistance, and quantity of electrical charge.
- Spectroscopic methods: Explore the interaction between electromagnetic radiation and analyte atoms or molecules or the emission of radiation by analytes.

What is role of analytical chemistry?



Fundamental Concepts

1- Atomic weight of element:

The mass of a single atom in grams is much too small a number for convenience, and chemists therefore use a unit called an *atomic mass unit (amu)* also known as a*dalton (Da)*. One *amu* is defined as exactly 1/12 the mass of carbon isotope 12 C and equal to 1 .66054 $\times 10^{-24}$ g.

Example: prove that carbon weighing 1.0×10^{-3} g contains 5.01×10^{19} carbon atom?

Solution:

2-Molecular weight.

The sum of the atomic weights of all the atoms in amolecule.

Molecular weight = Summation of atomic weight

Example: Calculate the molecular weight of water.

Solution:

A water molecule, H2O, has,

2 Hydrogen 2×1.0080 amu

1 Oxygen 1×15.9994 amu

Total molecular weight $= 18.0154 \text{ amu} \approx 18.02$

Example: Calculate the molecular weight of methyl alcohol.

Solution:

The molecular formula is CH₃OH or CH₄O. Then:

1 carbon $1 \times 12.011 \text{ amu} = 12.011 \text{ amu}$

4 hydrogen $4 \times 1.008 \text{ amu} = 4.032 \text{ amu}$

1 oxygen $1 \times 15.999 \text{ amu} = 15.999 \text{ amu}$

Total molecular weight = 32.04 amu

Home work.

Calculate the molecular weight of $C_{8}H_{18}$, $C_{14}H_{02}$, $C_{15}H_{15}SO_{24}$.

amu = 14.0067 for N₂ = 32.065 for S

3- Chemical equation.

- Representation of chemical reaction in terms of symbols and formula of reactants and products.
- Reactants written in left hand side.
- Products written in right hand side.
- numbers of atoms of different elements are the same on both sides of arrow is called equation. balanced chemical

Classification of chemical reactions.

1) Combination reaction:

$$2Mg(S) + O_2(g) \rightarrow 2MgO(s)$$

2) Decomposition reaction:

$$2Pb(NO_3)_2(s) \rightarrow 2PbO(s) + 4NO_2(g) + O_2(g)$$

3)Displacement reactions:

$$Zn(S) + CuSO_4(aq) \rightarrow ZnSO_4(aq) + Cu(s)$$

 $Cu(s) + 2AgNO_3(aq) \rightarrow Cu(NO_3)_2(aq) + 2Ag(s)$

4) Double Displacement reactions:

• Precipitation reaction:

$$AgNO_3(aq) + NaCl(aq) \rightarrow AgCl(s) + NaNO_3$$

• Neutralization reaction:

$$NaOH(aq) + HCl(aq) \rightarrow NaCl(aq) + H_2O$$

5)Redox reaction:

$$ZnO+C o Zn+CO$$
 ZnO reduce to Zn ------ Reduction O oxidize to CO ------ Oxidation

- **6)**Exothermic reaction and Endothermic reaction:
 - Endothermic reaction

$$CaCO_3 + Energy \rightarrow CaO + CO_2$$

• Exothermic reaction:

$$C + O_2 \rightarrow CO_2(g) + Energy$$

4-Mole (mol).

* Is the number of Avogadro's number of atom, molecule, electron and proton (6.022×10^{23}) .

The numbers of moles of substance is calculated from.

no. of moles =
$$\frac{mass\ of\ substance\ (g)}{molecular\ weight\ (\frac{g}{mol})}$$

 $Millimole (mmol) = 10^{-3} mol \ and \ 10^{3} mmol = 1 mol$

Example: Calculate the molar mass formaldehyde CH_2O .

Solution:

$$\begin{split} \mathcal{M}_{\text{CH}_2\text{O}} &= \frac{1 \text{ mol-C}}{\text{mol CH}_2\text{O}} \times \frac{12.0 \text{ g}}{\text{mol-C}} + \frac{2 \text{ mol-H}}{\text{mol CH}_2\text{O}} \times \frac{1.0 \text{ g}}{\text{mol-H}} \\ &+ \frac{1 \text{ mol-O}}{\text{mol CH}_2\text{O}} \times \frac{16.0 \text{ g}}{\text{mol-O}} \\ &= 30.0 \text{ g/mol CH}_2\text{O} \end{split}$$

Example: Find the number of moles and millimoles of benzoic acid (M = 122.1 g/mol) that are contained in 2.00 g of the pure acid.

Solution:

If we use HB_z to represent benzoic acid, we can write that 1 mole of HB_z has a mass of 122.1 g. Therefore,

amount HBz =
$$2.00 \text{ g HBz} \times \frac{1 \text{ mol HBz}}{122.1 \text{ g HBz}} = 0.0164 \text{ mol HBz}$$

To obtain the number of millimoles, we divide by the millimolar mass (0.1221 g/mmol), that is,

amount HBz =
$$2.00 \text{ g HBz} \times \frac{1 \text{ mmol HBz}}{0.1221 \text{ g HBz}} = 16.4 \text{ mmol HBz}$$

Or, no. of moles =
$$\frac{\text{mass of substance } (g)}{\text{molar mass } (\frac{g}{\text{mol}})}$$

no. of moles =
$$\frac{2 \text{ g}}{122.1(\frac{\text{g}}{\text{mol}})} = 0.0164 \text{ mol}$$

$$= 0.0164 \text{ mol} \times \frac{10^3 \text{ mmol}}{1 \text{ mol}} = 16.4 \text{ mmol HBz}$$

6-Equivalent weight.

Is the molecular weight divided by the number of reacting units (no.of equivalent = valency).

$$Equivalent Weight = molecular weight no. of equivalent$$

 For acid: the number of reacting units is the no. ofhydrogen ions that will furnish.

For example:

$$HCl \leftrightarrow H^+ + Cl^ eq.wt. HCl = \frac{M.wt.}{1}$$
 $H_2SO_4 \leftrightarrow 2H^+ + SO_4^{-2}$ $eq.wt. H_2SO_4 = \frac{M.wt.}{2}$ $CH_3COOH \leftrightarrow H^+ + CH_3COO^ eq.wt. CH_3COOH = \frac{M.wt.}{1}$

For Phosphoric Acid (H₃PO₄) the equivalent weight depends on the reaction as shown below.

$$\begin{array}{ll} H_3PO_4 \overset{M.O}{\Longleftrightarrow} H^+ + H_2PO_4^- & eq.wt. = \frac{M.wt.}{1} \\ H_3PO_4 \overset{Ph.ph}{\Longleftrightarrow} 2H^+ + HPO_4^{2-} & eq.wt. = \frac{M.wt.}{2} \\ H_3PO_4 \overset{CaCl_2}{\Longleftrightarrow} 3H^+ + PO_4^{3-} & eq.wt. = \frac{M.wt.}{3} \end{array}$$

> For bases: the number of reacting units is the no. of hydrogen ions that will react with it.

For example:

$$KOH + H^+ \to K^+ + H_2O$$
 $eq.wt. = M.wt.$
 $Ba(OH)_2 + 2H^+ \to Ba^{2+} + 2H_2O$ $eq.wt. = \frac{M.wt.}{2}$
 $Fe_3O_3 + 6H^+ \to 2Fe^{3+} + 3H_3O$ $eq.wt. = \frac{M.wt.}{6}$

> For salts:

$$eq.wt. = \frac{\textit{M.wt.of salt}}{\textit{no.of positive ion or no.of negative ion}}$$

For example:

$$Ag^{+}NO_{3}^{-} + H^{+}Cl^{-} \rightarrow Ag^{+}Cl^{-} + H^{+}NO_{3}^{-}$$
 $eq.wt. of Ag^{+}NO_{3}^{-} = \frac{M.wt. of Ag^{+}NO_{3}^{-}}{1}$
 $eq.wt. of Ag_{2}^{+}NO_{3}^{-2} = \frac{M.wt.}{2}$
 $eq.wt. of Al^{+3}PO_{4}^{-3} = \frac{M.wt.}{3}$
 $eq.wt. of Ca_{3}^{+2}(PO_{4})_{2}^{-3} = \frac{M.wt.}{6}$

> For oxidation / reduction reactions:

$$eq.wt. of \ reducing \ agent = \frac{\textit{M.wt.}}{\textit{no. of electrons loosed for one molcule}}$$

$$eq.wt. of \ oxidizing \ agent = \frac{\textit{M.wt.}}{\textit{no. of electrons accepted for one molcule}}$$

For example:

$$Mn^{+7}O_4^{-} + H^+ + 5e \rightarrow Mn^{+2} + H_2O$$
 In acidic medium eq. wt. of $Mn^{+7}O_4^{-} = \frac{M.wt.}{5}$

$$Mn^{+7}O_4^{-} + H_2O \rightarrow Mn^{+4}O_2^{-} + OH$$
 In basic medium eq. wt. of $Mn^{+7}O_4^{-} = \frac{M.wt.}{3}$

Home work:

Calculate the equivalent weights of the following substances as acids or bases: (a) HCl, (b) $Ba(OH)_2$, (c) $KH(IO_3)_2$, (d) H_2SO_3 , (e) CH_3COOH .