

Al-Mustaqbal University College
Department of Anesthesia techniques

Third Stage

Lecture 4

Asst. Lect. Raghda M. Alshemari

2021-2022





Frequencies and Distributions


- This is the most important univariate procedure.
- Conducting it properly, and interpreting the output rigorously, will enable you to understand the major attributes of the frequency distribution of each variable.



The distribution of variables – histograms and frequency statistics

- GO TO Analyze/ Descriptive Statistics/ Frequencies. Select the variables and move them into the box “Variable(s) as shown in Figure 1.”
- Unless the variables you have chosen are categorical or dummy (i.e. - they have only a few discrete possible values), deselect the option “Display Frequency Tables.” Otherwise, you will generate too many pages of output.

The distribution of variables – histograms and frequency statistics



Mean, Median, Mode, & Range

~~5~~, 7, 7, 9, ~~11~~, ~~12~~, 14

$$\text{Mean} = \frac{\text{Sum}}{n}$$

$$\text{Mode} = 7$$

$$\begin{aligned} \text{Range} &= 14 - 5 \\ &= 9 \end{aligned}$$


$$\text{Median} = 9$$



The distribution of variables – histograms and frequency statistics

- Conduct the frequencies procedure twice - Once for continuous variables (deselecting the option "Display Frequency Tables") and once for categorical and dummy variables (this time choosing the option "Display Frequency Tables").
- Now you must instruct SPSS to construct a histogram for each of the chosen variables. Click on the button "Charts."

The distribution of variables – histograms and frequency statistics






What is the normal distribution and why is it so important?


- A variable with a normal distribution has the same mode, mean, and median, i.e. - its most often occurring value equals the average of values and the mid-point of the values.
- Visually, a normal distribution is bell-shaped (as shown in Figure 3.) the "idealized normal curve- the left half is a mirror image of the right half.
- The importance stems from the assumption that "if a variable can be assumed to be distributed normally, then several inferences can be drawn easily and, more importantly, standardized tests (like the T and F tests) can be applied."

Histogram with normal distribution.



Applying descriptive by selecting statistics

- Click on the button “Statistics.” As in Figure 4. Select the options as shown in Figure 5. These statistics cover the list of "descriptive statistics.
- The options under “Percentile values” can assist in learning about the spread of the variable across its range.



Selecting required statistics

The screenshot shows the 'Frequencies: Statistics' dialog box with the following settings:

- Percentile Values:**
 - Quartiles
 - Cut points for: 10 equal groups
 - Percentile(s):
 - Buttons: Add, Change, Remove
- Central Tendency:**
 - Mean
 - Median
 - Mode
 - Sum
 - Values are group midpoints
- Dispersion:**
 - Std. deviation
 - Variance
 - Range
 - Minimum
 - Maximum
 - S.E. mean
- Distribution:**
 - Skewness
 - Kurtosis

Buttons at the bottom: Continue, Cancel, Help



Selecting required statistics

- In the next three graphs, the heights of the bars give the relative frequencies of the values of variables.
- Compare the bars (as a group) with the normal curve (drawn as a bell-shaped line curve). All three variables seem to be left heavy relative to the relevant normal curves, i.e. - lower values are observed more often than higher values for each of the variables, see Figure 6.
- On the lower-right corner, the chart provides the most important statistics - standard deviation, mean, and sample size. (The other statistics - like the median, mode, range, skewness, and kurtosis) are usually more visually identifiable from a histogram. The mode is the highest bar, the median has half the area (under the shaded bars) to its left, and the skewness and kurtosis are measures of attributes that are easily identifiable, see Figure 7.

Frequencies

→ Frequencies

Warnings

Treat is a string so a histogram cannot be produced.

Statistics

		Hospt	Treat
N	Valid	49	49
	Missing	0	0
Mean		2.08	
Median		2.00	
Mode		1	
Std. Deviation		1.152	
Variance		1.327	
Skewness		1.200	
Std. Error of Skewness		.340	
Kurtosis		1.122	
Std. Error of Kurtosis		.668	
Range		4	

Frequencies

Frequency Table

		Hospt			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1	18	36.7	36.7	36.7
	2	17	34.7	34.7	71.4
	3	10	20.4	20.4	91.8
	5	4	8.2	8.2	100.0
	Total	49	100.0	100.0	

		Treat			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Imipramine	18	36.7	36.7	36.7
	Lithium	18	36.7	36.7	73.5
	Placebo	13	26.5	26.5	100.0
	Total	49	100.0	100.0	



*Thank
You
For
Listening*