



Cell cycle

➤ Cell division in prokaryotes

- A. typically, a **prokaryotic cell divides by binary fission, splitting into two nearly equal halves**
- B. the main circular DNA molecule of the cell is replicated
- C. new plasma membrane and cell wall materials are laid down between the two DNA circles, eventually separating the daughter cells
- D. prokaryotic cells can have a **generation time (general term for the period from the start of one cell division to the start of the next cell division) as short as 20 minutes**

➤ Cell division in eukaryotic cell

Cell division is a very important process in all living organisms. The division cycle of most cells consists of four coordinated processes: **cell growth, DNA replication, distribution of the duplicated chromosomes to daughter cells, and cell division.**

Eukaryotic DNA molecules are organized in **chromosomes**

- each chromosome is made of **chromatin**, a long DNA molecule with associated proteins
- each chromosome contains hundreds to thousands of **genes**
- each species has a characteristic number of chromosomes
- chromosomes carry the genetic information of a cell from one cell generation to the next.
- Chromosome consists of two **chromatids or sister chromatids** join



Department of Anesthesia Techniques
Title of the lecture:- Cell cycle



Younis.Abdulridha@mustaqbal-college.edu.iq
Ataa_khalil@mustaqbal-college.edu.iq

together by **centromere**, which responsible for the movement of chromosomes at cell division.

Each species has a characteristic chromosome number; for instance, human cells contain 46 chromosomes, corn has 20 chromosomes.

- The full number or called **diploid (2n)** number of a chromosome that occurs in all cells of the body. The diploid number includes two chromosomes of each kind.
- Half the diploid number is called the **haploid (n)** number of chromosomes, representing only one of each kind of chromosome. In the life cycle of many animals, only sperm and eggs have the haploid number of chromosomes.

Cell cycle has two main phases interphase and cell division (mitosis + cytokinesis)

- **Interphase** is the longest stage in the eukaryote *cell cycle*. During interphase, the cell acquires nutrients, creates and uses proteins and other molecules, and starts the process of cell division by replicating the DNA. Interphase is divided into three distinct stages, *Gap 1*, *S*, and *Gap 2*. The purpose of interphase in all cell types is to **prepare for cell division, which happens in a different stage of the cell cycle.**

- **G₁ phase**

It is the post mitotic phase and first phase within interphase. After cells have finished dividing their chromosomes, and *cytokinesis* has divided the cell membrane, the two new cells enter the first stage of interphase, *Gap 1* or *G₁*. During this stage, the cell performs its



Department of Anesthesia Techniques
Title of the lecture:- Cell cycle



Younis.Abdulridha@mustaqbal-college.edu.iq
Ataa_khalil@mustaqbal-college.edu.iq

normal functions, and grows in size, the cell replicates organelles as necessary, cells double in organelles, accumulates materials needed for DNA synthesis and cell division

- **S phase**

It is the synthesis phase during this phase duplication of DNA and centriole takes places. The duplication of DNA results in the duplication of chromosomes. Thus, during this phase, the amount of DNA in the cell has effectively doubled

- **G2 phase**

It is the pre- mitotic phase (invisible phase) the synthesis of RNA and protein continues in this phase. The formation of macro molecules for spindle formation takes place and the cell prepare itself to go into the mitotic phase.

➤ **M phase or Mitotic phase**

During the mitotic (M) phase, the cell divides its copied DNA and cytoplasm to make two new cells. M phase involves: mitosis and cytokinesis.

1. **mitosis** is the process that distributes a complete copy of the duplicated genetic information to each daughter cell
 2. **cytokinesis** is the process of dividing the cytoplasm into two separate cells
- some cells can have mitosis without cytokinesis (most common in fungi and slime molds)

Mitosis is nuclear division that produces two daughter nuclei, each with the same number and kinds of chromosomes as the parental nucleus.



Department of Anesthesia Techniques
Title of the lecture:- Cell cycle



Younis.Abdulridha@mustaqbal-college.edu.iq
Ataa_khalil@mustaqbal-college.edu.iq

M phase is divided into four phases. These phases are; prophase, metaphase, anaphase, and telophase.

Prophase

During prophase, the chromosomes supercoil and the fibers of the mitotic spindle begin to form between centrosomes located at the pole of the cells. The nuclear membrane also disintegrates at this time, freeing the chromosomes into the surrounding cytoplasm.

Prometaphase

During prometaphase, some of the fibers attach to the centromere of each pair of sister chromatids and they begin to move toward the center of the cell.

Metaphase

At metaphase the chromosomes align along the center plane of the cell.

Anaphase

During anaphase, the centromeres split and the sister chromatids begin to migrate toward the opposite poles of the cell.

Telophase


During telophase, the chromosomes at either end of the cell begin to cluster together, which facilitates the formation of a new nuclear membrane. This also is when cytokinesis occurs, leading to two separate cells. One way to identify that telophase has begun is by looking for the formation of the cell plate, the new cell wall forming between the two cells.



Department of Anesthesia Techniques
 Title of the lecture:- Cell cycle



Younis.Abdulridha@mustaqbal-college.edu.iq
 Ataa_khalil@mustaqbal-college.edu.iq



Meiosis is a type of cell division that reduces the number of chromosomes in the parent cell by half and produces four gamete cells. This process is required to produce egg and sperm cells for sexual reproduction

Mitosis	Meiosis
1- Mitosis is the division of non sex cells 2- Create two identical daughter cells 3- With a complete set of chromosomes from both parents; 23 pairs chromosomes.	1- Meiosis is the division of sex cells 2- Create 4 different daughter cells, 3- With a half number of chromosomes each with <u>23 individual</u> chromosomes.