



كلية المستقبل الجامعة قُسم الفيزياء الطبية المرحلة الرابعة

# Medical Physics Neurophysics

## Lecture 4

Lecturer: Mohammed Salih

#### **Mechanism of Generated Action Potential :**

1- Stimulus starts the rapid change in voltage or action potential. In patch-clamp mode, sufficient current must be administered to the cell in order to raise the voltage above the threshold voltage to start membrane depolarization .

2- Depolarization is caused by a rapid rise in membrane potential opening of sodium channels in the cellular membrane, resulting in a large influx of sodium ions .

3- Membrane Repolarization results from rapid sodium channel inactivation as well as a large efflux of potassium ions resulting from activated potassium channels .



4- Hyperpolarization is a lowered membrane potential caused by the efflux of potassium ions and closing of the potassium channels .

5- Resting state is when membrane potential returns to the resting voltage that occurred before the stimulus occurred .

#### The Synapse :

Neurons essentially communicate with each other through synapses. When signals have traveled through neurons to the endpoint, they cannot simply continue onto the next neuron

Neurons do not touch each other, but where the neuron does come close to another neuron, a synapse is formed between the two .



The junction between the axon terminal of one neuron and the dendrite terminal of the other neuron is called a synapse. A small gap called the synaptic cleft separates it .

There are two types of synapses :

- 1-Electrical synapse.
- 2- Chemical synapse.

#### **Properties of synapse :**

Synapses also have the ability to communicate a change in the message being passed on. Postsynaptic neurons can also send communications back to the presynaptic neurons telling them to change how often or much a neurotransmitter is released.

Due to this, we can say that the synapses are able to communicate bidirectionally, A synapse is a combination of :

**1- Presynaptic endings** – which contain the neurotransmitters (chemical messengers ) .

2- Synaptic clefts – which is the gap between the two neurons .

**3- Postsynaptic endings** – which contains the sites for receptors (molecules which receives signals for a cell).



#### **Electrical and Chemical Synaptic Transmission?**

Synapses can be either chemical or electrical and are essential to the functioning of neural activity.

Neuroscientists understand that synapses play a vital role in a variety of cognitive functions, including learning and memory formation .

Electrical synapses are different from chemical synapses as there is a direct physical connection between the presynaptic and the postsynaptic neuron.

This connection takes the form of something called a gap junction, which is essentially a channel which allows ions to flow directly from the presynaptic cell to the postsynaptic cell .

Chemical Synapses	Electrical Synapses
1- Gap between cells is about 20	1- Gap between cells is about 3.5
nanometers	nanometers
2- Speed of transmission is several	2- Speed of transmission is nearly
milliseconds	instantaneous
3- Can be excitatory or inhibitory	3- Excitatory only
4- No loss of signal strength	4- Signal strength diminishes over
	time

### Signal Transmission Across a Synapse :

1- When an impulse reaches the far end (called the synaptic terminal), the impulse must travel from the presynaptic neuron to the postsynaptic neuron .

2- Chemical messengers called neurotransmitters carry the neural signal from one neuron to the next neuron or effector .



#### **Electrical Nature of Nerves :**

1- Neurons use electrical signals to communicate with other neurons, muscles and glands.

2- When microelectrodes are placed on either side of the membrane of an inactive neuron, measurements from a voltmeter indicate an electrical potential difference of -70mV (millivolts).

3- The charge of the inside of the neuron cell is negative in relation to the outside .This charge separation across the membrane is known as the membrane potential .