



2.2 Resultant of Coplanar Parallel Force System

Parallel forces can be in the same or in opposite directions. The magnitude of the parallel resultant force R is the magnitude of the algebraic sum of the given forces.



$$\uparrow^+ R = \sum F_i$$
$$R = -F_1 - F_2 + F_3 - F_4$$

The position of the resultant can be determined according to the principle of moments.

27

$$\bigoplus M_R = R \cdot d = \sum F_i \cdot d_i$$

 $R \cdot d = F_1 \cdot d_1 + F_2 \cdot d_2 - F_3 \cdot d_3 + F_4 \cdot d_4$





Dr.Mohammed Zuhair and Dr. Mayadah W.

Chapter Two: Resultant of Force Systems Resultant of Coplanar Parallel F. S.

Example No. 1: For the force system shown in figure, determine the magnitude and position of the resultant with respect to point A.



Solution:

 $M_{couple} = 165 \times 2 = 330 \ kN.m \ ' \ '$ $\rightarrow^{+} R = \sum F_{i}$ R = 200 + 50 - 250 - 150 $R = -150 \ kN = 150 \ kN \leftarrow$ $(+^{+} R \cdot d = \sum F_{i} \cdot d_{i} \quad (\text{Respect to } A)$ $-150 \cdot d = 200 \times 5 + 50 \times 2 - 250 \times 7 - 150 \times 4 + 330$ $d = \frac{-920}{-150}$ $d = 6.13 \ m$

> Dr.Mohammed Zuhair Dr. Mayadah W. Falah

Chapter Two: Resultant of Force Systems Resultant of Coplanar Parallel F. S.

Example No. 2: Find the value of P and F so that the four forces shown in Figure produce an upward resultant of 300 N acting at 4 m from point A.



Solution:

 $\uparrow^+ R = \sum F_i$ 200 N 100 N 300 = -100 + P - F + 2002 m 3 m 2 m $P = 200 + F \dots \dots \dots \dots (1)$ R = 300 N $d = 4^{\circ}m$ (+) $R \cdot d = \sum F_i \cdot d_i \quad (respect to A)$ $-300 \times 4 = 100 \times 0 - P \times 2 + F \times 5 - 200 \times 7$ $-2P + 5F - 200 = 0 \dots \dots \dots (2)$ Sub eq. (1) in eq. (2) to get: -2(200+F) + 5F - 200 = 0 $F = 200 N \downarrow$ Sub value of (F) in eq. (1) to get; $P = 400 N \uparrow$

Chapter Two: Resultant of Force Systems Resultant of Coplanar Parallel F. S.

Problems:

1. A parallel force system acts on the cantilever beam shown in Figure. Determine the magnitude and position of the resultant.



Answer: $R = 110 N \downarrow$, d = 6 m from point A

2. Compute the magnitude and position of the resultant of the two forces acting on a beam with respect to point A as shown in Figure.



Answer: $R = 80 N \leftarrow$, d = 40 mm up point A

3. The resultant of three parallel loads (one is missing in Figure below) is 13.6 kN acting up at 3 m to the right of A. Compute the magnitude and position of the missing load.



Answer: $F = 31.4 \ kN \downarrow$ at 2.48 *m* to the right of A