


# INTRODUCTION TO POLAR COORDINATES

- OBJECTIVES:** 1) Graph in polar coordinate form.  
2) Change from polar to rectangular coordinates.


## CARTESIAN VS. POLAR COORDINATES

Coordinate systems are used to describe the location of a point in space.

The Cartesian system describes how we should move from the origin both horizontally and vertically using coordinates  $(x,y)$ .






The polar system describes the distance straight from the origin to a point and determines the angle this segment makes with the positive x-axis.



In the polar system, we begin with a fixed point O in the plane called the **pole** (or **origin**) and draw from O a ray called the **polar axis**. Then each point P can be assigned polar coordinates  $P(r, \theta)$  where

- $r$  is the distance from O to P
- $\theta$  is the angle between the polar axis and the segment  $\overline{OP}$
- If  $\theta$  is positive, we measure counter-clockwise from polar axis. If  $\theta$  is negative, we measure in clockwise direction.
- If  $r$  is negative, then  $P(r, \theta)$  is  $r$  units from the pole in the opposite direction of  $\theta$ .



- Plot the following points:

$$A\left(1, \frac{3\pi}{4}\right) \quad B\left(3, -\frac{\pi}{6}\right) \quad C(3, 3\pi) \quad D\left(-4, \frac{\pi}{4}\right)$$


- $P(r, \theta)$  can be represented by:

$$P(r, \theta + 2\pi n) \quad \text{or} \quad P(-r, \theta + (2n + 1)\pi)$$

- If  $P\left(2, \frac{\pi}{3}\right)$ , list four other polar coordinates for P.

$$\begin{array}{cc} r > 0 & r < 0 \\ \left(2, \frac{7\pi}{3}\right) \left(2, -\frac{5\pi}{3}\right) & (-2, \frac{4\pi}{3}) (-2, -\frac{2\pi}{3}) \end{array}$$

## CONVERTING POLAR/RECTANGULAR COORDINATES



From the diagram, we arrive at the following relationships using Pythagorean Theorem and right-triangle trig:

$$x^2 + y^2 = r^2$$

$$\cos \theta = \frac{x}{r} \quad \sin \theta = \frac{y}{r} \quad \tan \theta = \frac{y}{x}$$

$$r \cos \theta = x \quad r \sin \theta = y$$

1) Convert each rectangular point to polar coordinates:

a)  $(-1, 1)$   $x^2 + y^2 = r^2$   
 $r = \pm \sqrt{2}$

$$\tan \theta = \frac{y}{x} \quad \tan \theta = -1$$

$$\theta = \frac{3\pi}{4} \text{ or } -\frac{\pi}{4}$$

$(-1, 1)$  lies in quad 2!  $(\sqrt{2}, \frac{3\pi}{4})$  or  $(-\sqrt{2}, -\frac{\pi}{4})$

b)  $(0, 2)$   $x^2 + y^2 = r^2$   
 $r = \pm 2$

$$\tan \theta = \frac{2}{0} \leftarrow \text{undefined!}$$

$$\theta = \frac{\pi}{2} \text{ or } -\frac{\pi}{2} \quad \left(2, \frac{\pi}{2}\right)$$

or  
 $(-2, -\frac{\pi}{2})$

2) Given the polar coordinates, find the rectangular coordinates

a)  $\left(5, \frac{2\pi}{3}\right)$   $x = r \cos \theta$   $y = r \sin \theta$

$$x = 5 \cos\left(\frac{2\pi}{3}\right) \quad y = 5 \sin\left(\frac{2\pi}{3}\right)$$

$$x = 5\left(-\frac{1}{2}\right) \quad y = 5\left(\frac{\sqrt{3}}{2}\right)$$

$$x = -\frac{5}{2} \quad y = \frac{5\sqrt{3}}{2} \quad \left(-\frac{5}{2}, \frac{5\sqrt{3}}{2}\right)$$

b)  $\left(\sqrt{3}, \frac{\pi}{6}\right)$

$$x = \sqrt{3} \cos\left(\frac{\pi}{6}\right) \quad y = \sqrt{3} \sin\left(\frac{\pi}{6}\right)$$

$$x = \sqrt{3} \cdot \frac{\sqrt{3}}{2} \quad y = \sqrt{3} \cdot \frac{1}{2}$$

$$x = \frac{3}{2} \quad y = \frac{\sqrt{3}}{2} \quad \left(\frac{3}{2}, \frac{\sqrt{3}}{2}\right)$$

## CONVERTING A RECTANGULAR EQUATION TO A POLAR FORM

3) Convert the rectangular equation to polar form:  $x^2 = 4y$

$$(r \cos \theta)^2 = 4r \sin \theta$$

$$r^2 \cos^2 \theta = 4r \sin \theta$$

$$r = \frac{4r \sin \theta}{r \cos^2 \theta}$$

$$r = \frac{4 \tan \theta}{\cos \theta}$$

$$\boxed{r = 4 \tan \theta \sec \theta}$$

Use direct conversion

4) Convert the rectangular equation to polar form:  $4x + 7y - 2 = 0$

$$4r \cos \theta + 7r \sin \theta - 2 = 0$$

$$r(4 \cos \theta + 7 \sin \theta) = 2$$

$$\boxed{r = \frac{2}{4 \cos \theta + 7 \sin \theta}}$$

5) Convert the rectangular equation to polar form:  $x^2 + y^2 - 8y = 0$

$$r^2 - 8r \sin \theta = 0$$

$$r(r - 8 \sin \theta) = 0$$

$$r - 8 \sin \theta = \frac{0}{r}$$

$$\boxed{r = 8 \sin \theta}$$

notice patterns:  $x^2 + y^2 = r^2$

## CONVERTING A POLAR EQUATION TO RECTANGULAR FORM


Convert the polar equation to rectangular form. If possible, determine the graph of the equation from its rectangular form.

6)  $r = 5 \sec \theta$

$$r = \frac{5}{\cos \theta}$$

$$r \cos \theta = 5$$

$$x = 5$$



7)  $r = 2 \sin \theta$

$$r = 2 \sin \theta$$

$$r^2 = 2r \sin \theta$$


$$x^2 + y^2 = 2y$$

$$x^2 + y^2 - 2y = 0$$

$$x^2 + y^2 - 2y + 1 = 1$$

$$x^2 + (y-1)^2 = 1$$

Circle: center at  $(0,1)$   
radius 1



8)  $r = 2 + 2 \cos \theta$

$$r^2 = 2r + 2r \cos \theta$$

$$x^2 + y^2 = 2r + 2x$$

$$(x^2 + y^2 - 2x)^2 = (2r)^2$$

$$(x^2 + y^2 - 2x)^2 = 4r^2$$

$$(x^2 + y^2 - 2x)^2 = 4(x^2 + y^2)$$

Yuck!

9)  $r = \sin\left(\theta - \frac{\pi}{4}\right)$

$$r = \sin \theta \cos\left(\frac{\pi}{4}\right) - \cos \theta \sin\left(\frac{\pi}{4}\right)$$

$$r = \frac{\sqrt{2}}{2} \sin \theta - \frac{\sqrt{2}}{2} \cos \theta$$

$$r^2 = \frac{\sqrt{2}}{2} r \sin \theta - \frac{\sqrt{2}}{2} r \cos \theta$$

$$x^2 + y^2 = \frac{\sqrt{2}}{2} y - \frac{\sqrt{2}}{2} x$$


$$x^2 + \frac{\sqrt{2}}{2} x + y^2 - \frac{\sqrt{2}}{2} y = 0$$

$$x^2 + \frac{\sqrt{2}}{2} x + \frac{1}{8} + y^2 - \frac{\sqrt{2}}{2} y + \frac{1}{8} = \frac{1}{8} + \frac{1}{8}$$

$$\left(x + \frac{\sqrt{2}}{4}\right)^2 + \left(y - \frac{\sqrt{2}}{4}\right)^2 = \frac{1}{4}$$

### THE POLAR DISTANCE FORMULA:

$$d^2 = r_1^2 + r_2^2 - 2r_1r_2 \cos(\theta_1 - \theta_2)$$



10) Find the distance from  $\left(2, \frac{\pi}{6}\right)$  to  $\left(-3, \frac{\pi}{4}\right)$ .


$$d^2 = 2^2 + (-3)^2 - 2(2)(-3)\cos\left(\frac{\pi}{6} - \frac{\pi}{4}\right)$$

$$d = \sqrt{4 + 9 + 12\cos\left(-\frac{\pi}{12}\right)}$$

### THE POLAR CIRCLE FORMULA:

$$a^2 = r^2 + r_0^2 - 2rr_0 \cos(\theta - \theta_0)$$

Radius  $a$  and center  $(r_0, \theta_0)$



Find the equation of a circle with:

11) radius 5 and center  $(0, 0)$  <sup>origin</sup>

$$r = 5$$

12) radius 1 and center  $\left(2, \frac{\pi}{6}\right)$

$$1 = r^2 + 4 - 2(r)(2)\cos\left(\theta - \frac{\pi}{6}\right)$$

$$1 = r^2 + 4 - 4r\cos\left(\theta - \frac{\pi}{6}\right)$$