

Lec5

Inhalational anesthetics

Osama Aziz & Muaid abdallah B.M.TCH
ANAESTHESIA
2023 -2024

Al_Mustaqbal University

College of Health and Medical Techniques Anesthesia Techniques
Department



Definition: An inhalational anesthetic is a chemical compound possessing general anesthetic properties that can be delivered via inhalation. They are administered through a face mask, laryngeal mask airway or tracheal tube connected to an anesthetic vaporizer and an anesthetic delivery system.

The main use of inhalational was to maintain anesthesia, some of them can be used as induction

The dose of inhalational anesthetic was mentioned as MAC

MAC: The minimum alveolar concentration (MAC) of an inhaled anesthetic is the alveolar concentration that prevents movement in 50% of patients in response to a standardized stimulus (eg, surgical incision). e.g : Halothane MAC 0.75%

Factors Increasing MAC (need more volatile to maintain anesthesia)

- Hyperthermia
- Hyponatraemia
- Sympatho-adrenal stimulation
- Chronic alcohol abuse
- Chronic opioid abuse
- Increases in ambient pressure
- Hypercapnia
- Decreasing age
- Thyrotoxicosis

Factors decreasing MAC (need less volatile to maintain anesthesia)


- Nitrous oxide
- Hypothyroid/myxedema
- Hypocapnia
- Hypothermia-decrease is roughly linear
- Hyponatraemia
- Increasing age
- Hypoxaemia
- Hypotension
- Anemia

- Pregnancy
- CNS depressant drugs
 - Other drugs: lithium, lidocaine, magnesium, Acute alcohol abuse

HINT: Sex, Weight and Duration of anesthesia does not affect MAC
Nitrous oxide MAC 105% , Halothane (Fluothane) 0.75% , Isoflurane 1.2% , Desflurane 6.0% , Sevoflurane 2.0 %

Characteristic of IDEAL inhalational:

- Non-flammable, non-explosive at room temperature
- Stable in light.
- Liquid and vaporizable at room temperature
- Stable at room temperature, with a long shelf life
- Stable with soda lime, as well as plastics and metals
- Environmentally friendly - no ozone depletion
- Cheap and easy to manufacture
- Non toxic
- Rapid induction and rapid recovery



FGF (fresh gas flow) is determined by the vaporizer and flowmeter settings.

F_i (inspired gas concentration) is determined by (1) FGF rate; (2) breathing-circuit volume; and (3) circuit absorption.

F_A (alveolar gas concentration) is determined by (1) uptake (uptake = $\lambda \cdot b/g \times C(A-V) \times Q$); (2) ventilation; and (3) the concentration effect and second gas effect:
 a) concentrating effect
 b) augmented inflow effect

F_a (arterial gas concentration) is affected by ventilation/perfusion

- Safe with no toxic side effect
- There was no ideal inhalation till now

Nitrous oxide (N₂O): Inhalational anesthetic agent, being a useful analgesic for dental extraction



Properties:

- Colorless
- Slightly sweet-smelling gas
- MAC 105%.
- Non-flammable but supports combustion
- Breaking down to O₂ and nitrogen at high temperatures.

- Supplied as a liquid/gas in French blue cylinders
- Ice often forms on the cylinder during use

CNS effect:

- Fast onset and recovery; strongly analgesic but weakly anaesthetic.

Increases cerebral metabolism, cerebral blood flow and ICP slightly.

Respiratory System:

- Non-irritant. Depresses respiration slightly.
- May cause diffusion hypoxia at the end of surgery.
- Cardiovascular system:
- Little effect on heart rate and BP usually

Other:

- Post operative nausea and vomiting

- Does not affect hepatic or renal function, nor uterine or skeletal muscle tone.
- Prolonged use may cause bone marrow depression, megaloblastic anaemia and peripheral neuropathy.
- Generally considered as being safe during pregnancy

Halothane: Halothane is a halogenated alkane, it was nonflammable and non-explosive. Sensitive to light so thymol preservative and amber-colored bottles retard spontaneous oxidative decomposition. Its use rapidly spread because of its greater potency, ease of use, non-irritability and noninflammability, Risks of arrhythmias and

liver damage on repeated administration
(halothane hepatitis)

Properties:

- Colorless liquid
- Pleasant smell
- MAC 0.75%.
- Non-flammable.
- Supplied in liquid form with thymol 0.01%
- Decomposes slightly in light.

Central nervous system Effects:

- Smooth rapid induction, with rapid recovery.
- Anticonvulsant action.
- Increases cerebral blood flow but reduces intraocular pressure.



Respiratory system:

- Non-irritant. Pharyngeal, laryngeal and cough reflexes are abolished early
- Respiratory depressant, with increased respiratory rate and reduced tidal volume.

- Bronchodilatation and inhibition of secretions.

Cardiovascular system:

- Myocardial depression and bradycardia.
- Hypotension is common.
- Myocardial O₂ demand decreases.
- Arrhythmias are common, e.g.

Bradycardia ,ectopic

- Sensitizes the myocardium to catecholamines, e.g. Endogenous or injected adrenaline.

Other:

- Dose-dependent uterine relaxation.
- Nausea/vomiting is uncommon.
- May precipitate Malignant Hyperthermia.
- Up to 20% is metabolized in the liver.

- Repeat administration after recent use may result in hepatitis.

Isoflurane:



Properties:

- Colorless liquid
- Pungent odor
- MAC 1.20
- Non-flammable, non-corrosive.
- With no additive.

Central nervous system Effects:

- Smooth, rapid induction, but speed of uptake is limited by respiratory irritation.
- Recovery is slower than with sevoflurane and desflurane.
- Anticonvulsant properties
- Reduces Cerebral Metabolic Rate of O₂.
- Increases cerebral blood flow and ICP.

- Decreases intraocular pressure.
- Has poor analgesic properties.
- Respiratory System:
 - Irritant; more likely to cause coughing and laryngospasm
 - Respiratory depressant, with increased rate and decreased tidal volume.
 - Causes bronchodilatation.

- Cardiovascular system:
 - Myocardial depression is less than with halothane
 - Vasodilatation and hypotension commonly occur
 - Compensatory tachycardia is common
 - Myocardial O₂ demand decreases, but tachycardia may reduce
 - myocardial O₂ supply.

- • Coronary steal may occur
- Other:
 - • Dose-dependent uterine relaxation.
 - • Nausea/vomiting is uncommon.
 - • Skeletal muscle relaxation
 - • May precipitate MH.
 - • Widely used in neurosurgery

Sevoflurane



Properties:

- Colorless liquid
- Pleasant smelling
- MAC2.
- Non-flammable, non-corrosive.
- Supplied in liquid form with no additive.

- Interacts with soda lime to produce compounds A

Central nervous system Effects:

- Smooth, extremely rapid induction and recovery.
- Early postoperative analgesia may be required as emergence
- Is so rapid.

- Increases the risk of emergence agitation
- Anticonvulsant properties.
- Reduces cerebral metabolic rate of O₂
- Decreases intraocular pressure.
- Has poor analgesic properties.

Respiratory System:

- Well-tolerated
- Minimal airway irritation.

- Respiratory depressant, with increased rate and decreased tidal volume.
- Causes bronchodilatation.

Cardiovascular system:

- Vasodilatation and hypotension may occur
- Myocardial O₂ demand decreases.
- Arrhythmias uncommon

Other:

- Dose-dependent uterine relaxation.
- Nausea/vomiting occurs.
- Skeletal muscle relaxation
- May precipitate MH.
- Tracheal intubation may be performed easily with spontaneous Respiration.

- Considered the agent of choice for inhalational induction in pediatrics because of its rapid and smooth induction characteristics.
- Has also been used for the difficult airway, including airway obstruction.

Desflurane

Introduced in the UK in 1994, a colorless liquid with slightly pungent vapor, boiling point: 23°C, MAC: 5%–7% in adults; 7.2%–10.7% in children, non-flammable, non-corrosive, supplied in liquid form with no additive, may react with dry soda lime to produce carbon monoxide, requires the use of an electrically powered vaporizer due to its low boiling point.

Effects:

On central nervous system:

- 1) Rapid induction (although limited by its irritant properties) and recovery.
- 2) May increase cerebral blood flow, although the response of cerebral vessels to CO₂ is preserved.

3) ICP may increase due to imbalance between the production and absorption of CSF.

4) Reduces CMRO₂ as for isoflurane.

5) Has poor analgesic properties.

On respiratory system:

1) Causes airway irritation; not recommended for induction of anesthesia because respiratory complications (e.g. laryngospasm, breath-holding, cough, apnea) are common and may be severe.

2) Respiratory depressant, with increased rate and decreased tidal volume.

On cardiovascular system:

1) Vasodilatation and hypotension may occur, similar to isoflurane, may cause tachycardia and hypertension via sympathetic stimulation, especially if high concentrations are introduced rapidly.

2) Myocardial ischemia may occur if sympathetic stimulation is excessive, but has cardioprotective effects in patients undergoing cardiac surgery.

3) Arrhythmia as uncommon, as for isoflurane, little myocardial sensitization to catecholamines.

4) Renal and hepatic blood flow generally preserved.

Others:

1) Dose-dependent uterine relaxation (although less than isoflurane and sevoflurane).

2) Skeletal muscle relaxation; non-depolarising neuromuscular blockade may be potentiated.