

Lecture-3-

Physiology of Bacteria:

Dr.Nada Khazal K. Hendi

# L3: Physiology of Bacteria:

## Bacterial growth

**Growth** is the orderly increase in the sum of all the components of an organism. **Cell multiplication** is a consequence of growth, in unicellular organism, growth leads to an increase in the number of individuals making up a population. The bacterial growth can be measured by:

A: cell concentration: viable cell count.

B: Bio mass density: by determining the dry weight of a microbial culture.

## Phases of the bacteria growth:

The phases of the bacterial growth curve are shown in figure:

- 1\_lag phase:** represents the period of adaptation to the new environment, enzymes. And intermediates are formed.
- 2\_log (exponential) phase:** the number of cell increase exponentially (one becomes 2,  $2 \rightarrow 4$ ,  $4 \rightarrow 8$ .....) , the time required for this doubling is called as generation time or doubling time.
- 3\_stationary phase:** the exhaustion of nutrients or the accumulation of toxic products
- 4\_death phase:** after the period of time in the stationary phase, and completely exhausted of nutrients and accumulation of high level of toxic substances, the death increase until it reach a steady level.

**Table (3) Phase & Growth rate of the bacterial growth**

<b>Phase</b>	<b>Growth rate</b>
<b>Lag</b>	<b>Zero</b>
<b>Exponential</b>	<b>Positive constant</b>
<b>Stationary</b>	<b>Zero</b>
<b>Death</b>	<b>Negative (death)</b>




Figure (3) the phases of the bacterial growth curve

# Environmental growth factors

## A-Nutrients

The following nutrients must be provided for any bacterial culture:

1. Hydrogen donors and accepters.
2. Carbon source.
3. Minerals elements (sulfur and phosphorus).
4. Growth factors (amino acid, pyrimidines and purin).

## B-pH:

bacteria can be classified according to pH rang:

1. Neutrophiles: pH=6-8. most pathogenic bacteria.
2. Acidophiles: pH=3. can grow at pH as low as 3.
3. Alkaliphiles: pH=10.5. can grow at pH as high as 10.5.

C- Temperature: bacteria can be classified according to temperature:

1. Mesophiles: grow best at 30-37°C. most pathogenic bacteria.
2. Psychophiles: grow best at 15-20°C.
3. Thermophiles: grow best at 50-60°C.

D- Oxygen:

1. Obligate aerobes: need O<sub>2</sub> as hydrogen acceptor.
2. Obligate anaerobes:- need substance other than O<sub>2</sub> , and being sensitive to O<sub>2</sub> inhibition
3. Facultative:- able to live aerobically or an aerobically.

E-salt & osmotic pressure

- ♣ Halophiles: requiring high salt concentrations (marine bacteria).
- ♣ Osmophiles: requiring high osmotic pressure.

**aerobic bacteria** +O<sub>2</sub> → H<sub>2</sub>O<sub>2</sub> (toxic for bacteria ) →produce enzyme by bacteria (catalase) → H<sub>2</sub>O+ O<sub>2</sub> (gas bubbles)

**Non aerobic bacteria** +O<sub>2</sub> → H<sub>2</sub>O<sub>2</sub> (toxic for bacteria ) →bacteria is not produce enzyme) → H<sub>2</sub>O<sub>2</sub>



# Microbiological culture media

- The growth of MO depends on available nutrients and favorable growth environment. The cultures media differ depend on MO needing to nutrition. The media divided depending on their **contents** into:

1. Natural media.

2. Synthetic media.

3. Semi synthetic media.

- While the media can be divided **depending physical state** in to:

1. Liquid media (broth).

2. Solid media (agar).

3. Semisolid media.

- Type of inoculation of media:

1. Streaking plate. 2. Spreading methods.

3. Pour plate technique. 4. Stapping methods.

## stains:

are chemical compounds with colored ions which react with the different constituents of the living cell to stain the transparent and minute cells of bacteria which are difficult to see by naked eyes.

The stain may be classified into simple and differential, the simple stain colored all parts of the cell e.g. methylene blue while the differential stain are so selected that react with specific groups of the different parts of the cell or of different genera and species of bacteria for example Gram stain helps in the identification of bacteria G+ ve and G-ve especially the pathogenic agents that causing disease in the lab. Many theories have been proposed to explain the observed difference in Gram staining, one of the most common theory is that based on variation in the chemical composition of bacterial cell wall . G+ve bacteria contain magnesium- RNA-protein – carbohydrate complex which form an insoluble substance with the crystal violet and iodine , this complex is not washed with alcohol , while the lipid content of the cell wall being 10 times in G-ve as much as in G+ve ones , this lead to the solubility of lipids in alcohol and increase of cell wall porosity in G-ve and crystal violet iodine –complex can be extracted , so , G-ve cells become colorless after alcohol washing and take the another color safranin (cell turn pink) as shown in the below figure .

# Gram stain

G+ ve

G-ve

**-crystal violet**

Washing after 1 mint

purple

purple

**-Iodine**

Washing after 1 mint

purple

purple

**-alcohol**

Washing after 15 sec

purple

colorless

**-Safranin**

Washing after 1 mint

purple

pink



# Source of metabolic energy:

The three major mechanisms for generating metabolic energy are fermentation, aerobic respiration, and anaerobic respiration:-

1. Aerobic respiration:- is the metabolic process in which ( $O_2$ ) serves as the final electron acceptor for the electron transport chain. In this process  $O_2$  is reduced to water. This is the energy-generating used by all aerobic bacteria.
2. Anaerobic respiration: - is the metabolic process in which inorganic compounds **rather than  $O_2$**  serve as the final acceptors, these acceptors can be nitrate or sulfate.

3. **Fermentation:** is an alternative aerobic process, by which an organic metabolic intermediate serves as the final electron transport, such as **pyruvate, lactate, or ethanol** which is released from glucose fermentation.
- In all pathways the energy represented by ATP is released and used in the biosynthesis of bacterial cell component (cell wall, capsule,...) the ATP yielded is much higher in respiration than fermentation.