



Al-Mustaqbal University / Nursing College
Academic Year 2023-2024
Epidemiology



Lecture 4

Measurements in Epidemiology

By

Dr. Ghassan Abdul-Ameer

- **The four types of epidemiologic measures are:**

- 1. Count
- 2. Ratio
- 3. Proportion
- 4. Rate

Count

- ❑ The simplest and most frequently performed quantitative measurement in epidemiology is **a count**.
- ❑ Count of the number of persons in the group who have a particular disease or a particular characteristics.
- ❑ E.g. cases of influenza reported in New York, during January of a particular year.

Ratio

- **Ratio** is numerical expression, which indicates the relationship in quantity between two parts.
- Obtained by dividing one quantity by another.
- Example: (20) male patients were attended the clinic and (10) female patients the ratio of male to female is 2:1.

Of 1,000 motorcycle fatalities, 950 victims are men and 50 are women. The sex ratio for motorcycle fatalities is:

$$\frac{\text{Number of male cases}}{\text{Number of female cases}} = \frac{950}{50} = 19:1 \text{ male to female}$$

Proportion:

For a count to be descriptive of a group it must be seen in proportion to it; that is must be divided by the total number in the group.

- proportions may be expressed as percentages (%)

Calculation of the Proportion of African-American Male Deaths Among African-American and White Boys Aged 5 to 14 Years

A	B	Total (A + B)
Number of deaths among African-American boys 1,150	Number of deaths among white boys 3,810	Total 4,960

$$\text{Proportion} = A/(A + B) \times 100 = (1,150/4,960) \times 100 = 23.2\%$$

Rate

- **Rate**: is the quantity amount or degree of something measured in **specific period of time**.
- It is similar to proportion but it is calculated in a specific period(**usually one year**)

- **Epidemiologic rates contain the following elements:**
- 1- Disease frequency.
- 2- Unit size of population.
- 3- Time period during which an event occurs.

The crude death rate: reflects all deaths in the population regardless of age or cause of death. The crude death rate presents a picture of the overall health status of the population.

(e.g. all cancer deaths in 2000).

Number of death

Crude Death Rate = ----- * 1000

Number of deaths Estimated midinterval population

Specific Rate: Rates for specific segments/groups of the population (e.g. sex, age, race, cause of death, cancer site).

$$\textit{Age – specific death rate} = \frac{\textit{Number of deaths of a specified age group}}{\textit{Estimated midinterval population of that age group}} \times 1000$$

Prevalence

- The number of existing cases depends on the number of people who developed their illness in the past and have continued to be ill at the present time (i.e. **old** and **new**).

$$\textit{Prevalence} = \frac{\textit{Total Number of (old + new) cases over a time period}}{\textit{Total population at risk during the same time period}} \times (\textit{e.g., 100,000})$$

Incidence

- Is defined as the number (No.) of new cases of disease.
- **Incidence** is a measure of a disease developed in a person who did not have the disease.
- Transition from non disease to disease status.

Incidence rate

- No. of new cases in the population during specific period/ no of population at risk of developing the disease at that period.

Incidence is the measure of **disease developing** in a person who did not have the disease .

$$\text{Incidence rate} = \frac{\text{Number of new cases over a time period}}{\text{Total population at risk during the same time period}} \times \text{Multiplier (e.g., 100,000)}$$

Factors of Influencing prevalence rate :

- 1.The number of new cases(incidence):** if the incidence increase the prevalence will increase.
- 2.The severity of illness :**if the deaths increase the prevalence will decrease ,if the disease is easily cured the prevalence also decrease.
- 3.The duration of illness:** if the duration of the disease increase the prevalence will increase.


Period prevalence:

- The prevalence of a particular disease in a longer period such as one month , six months , one year or even more (life time prevalence).

Point prevalence :

- Is used to know how much of a particular disease is present in a population at a particular point in time(one day) ,
 - example Dec.31,2019.

Relationship between incidence and prevalence



Vital statistics:

- Tools used to **evaluate** the **health** status of the **community** .
- By vital statistics we refer to data collected from ongoing recording ,registration of all vital events (birth, deaths ,fetal deaths ,marriages and divorces).
- The nurse or physician is responsible for filling the certificates of death , birth and fetal death

Crude Birth Rate (CBR)

Crude Birth Rate (CBR) : Is the No. of live births per 1000 population of certain locality and year .

Number of live birth (during a year)

-----X (1000)

Mid year population in certain locality

**Birth rate is generally high in developing countries including Iraq ,
due to :**

1- high fertility rates due social and traditional motives .

2- poor family planning facilities

**3– factors related to marriage (the youngest age of marriage is
associated with long childbearing period .**

Crude Death Rate (CDR) =

$$\frac{\text{Total number of deaths for all ages and causes}}{\text{Mid year population (MYP)}} * 1000$$

$$\text{Neonatal Mortality Rate} = \frac{\text{Number of deaths under 28 days during a given time interval}}{\text{Number of live births during the same time interval}} \times 1000$$

$$\text{Infant Mortality Rate} = \frac{\text{Number of deaths under one year of age during a given time interval}}{\text{Number of live births during the same time interval}} \times 1000$$

$$\text{Maternal Mortality Rate} = \frac{\text{Number of deaths assigned to pregnancy related causes during a given time interval}}{\text{Number of live births during the same time interval}} \times 1000$$

Thanks