Department of Radiology Techniques Radiological Position The Second Stage



Forearm And Types of Fractures Lecture 4

Assist. Lecturer Tariq Nadhim Jassim

Basic Projections of Forearm

Two Positions

1- Anterior - Posterior (AP)

2- Lateral

Cassette Out Bucky .

24 x 30-cm (10x12 inch) Cassette Size



Styloid Process ———————————————————————————————————	wik Styloid Process
Ulna ————————————————————————————————————	Radius
Trochlea ————	——— Radial Tubercle ——— Radial Neck ——— Radial Head ——— Capitulum
Dlecranon Process ———	——— Humerus

1- Anterior - Posterior (AP)

Position of Patient

- The patient is seated alongside the table, with the affected side nearest to the table.
- The arm is abducted and the elbow joint is fully extended, with the supinated forearm resting on the table.
- The shoulder is lowered to the same level as the elbow joint.
- The arm is adjusted such that the radial and ulnar styloid processes and the medial and lateral epicondyles are equidistant from the cassette.
- The lower end of the humerus and the hand are immobilized using sandbags

Direction and centering of the X-ray beam


• The vertical central ray is centered in **the midline of the forearm to a point midway between the wrist** and elbow joints.


Essential image characteristics

 Both joints elbow and the wrist should be seen in the true anterio-posterior position, with the radial and ulnar styloid processes and the epicondyles of the humerus equidistant from the cassette.

Note

The posterior-anterior projection of the forearm with the wrist pronated is not satisfactory because, in this projection, the radius is superimposed over the ulna for part of its length





2- Lateral

Position of Patient

- From the anterior-posterior position, the elbow is flexed to 90 degrees.
- The humerus is internally rotated to 90 degrees to bring the medial aspect of the upper arm, elbow, forearm, wrist and hand into contact with the table.
- The cassette is placed under the forearm to include the wrist joint and the elbow joint.
- The arm is adjusted such that the radial and ulnar styloid processes and the medial and lateral epicondyles are superimposed.
- The lower end of the humerus and the hand are immobilized using sandbags.

Direction and Centering of the X-ray beam

• The vertical central ray is centered in **the midline of the forearm to a point midway between the wrist** and elbow joints.

Essential Image Characteristics

- Both the elbow and the wrist joint must be demonstrated on the image.
- Both joints should be seen in the true lateral position, with the radial and ulnar styloid processes and the epicondyles of the humerus superimposed.





Bone Fracture

A bone fracture is a full or partial break in the continuity of bone tissue. Fractures can occur in any bone in the body.

Types of Bone Fractures divided according to

Displaced Fracture: bone breaks into two or more pieces and moves out of alignment.

Non-Displaced Fracture: the bone breaks but does not move out of alignment (e.g.

greenstick fracture)

Closed Fracture: the skin is not broken.

Open Fracture: the bone has broken through the skin(compound or complex) this is a **medical**


emergency. In Addition to

- **1-Avulsion Fracture:** when a fragment of bone is pulled away by ligament or tendon which its attachment.
- **2-Impacted Fracture:** ends are driven into each other; commonly seen in arm fractures in children.
- **3-Comminuted Fracture:** the bone breaks into several pieces.
- 4-Compression or Wedge Fracture: usually involves the bones in the back (vertebrae).
- **5-Greenstick Fracture:** an incomplete fracture in which the bone is bent; occurs most often in children.
- **6-Linear Fracture:** the break is parallel to the bone's long axis.
- **7-Oblique Fracture:** the break has a curved or sloped pattern.

- **8-Pathologic Fracture:** caused by a disease that weakens the bones.
- 9-Spiral Fracture: one part of the bone has been twisted at the break point.
- 10-Stress Fracture: a hairline crack.
- 11-Transverse Fracture: the broken piece of bone is at a right angle to the bone's axis.
- 12-Epiphyseal Fracture: a fracture through the epiphysis.
- 13-Depressed Fracture: a fracture in which fragments are driven inward (skull and facial fracture)
- 14-Segmental Fracture: The same bone is fractured in two places, segment of bone between the

two breaks.

TYPES OF BONE FRACTURES





External Fixation Internal Fixation Oblique Fracture Avulsion Fracture



Wedge Fracture

Pathologic Fracture

Greenstick Fracture

