



## Vectors, Vectors in Space, Unit Vector, Scalar Product, Vector Product

### المتجه

#### Vectors و المتجهات


The vector is only two pieces of information:

- 1- Direction
- 2- Length or Magnitude

المتجه عبارة عن مكونين وهما : اتجاه و طول أو كبير.

We can graph a vector by an arrow that we can visualize on x-y plane and we can capture it by the arrow length and angle

Vectors on graph could start from not just an origin, but from anywhere.



"Examples of vectors"

#### Examples of vectors و المتجهات

To answer the question "What is the current temperature?" we use a single number (scalar) ; likewise the question about a mass ;

While to answer the question "What is the current velocity of the wind?" , we need more than just a single number. We need magnitude (speed) and direction. This where vectors come to handy.

Position, displacement, velocity, acceleration, force, momentum & torque are all physical quantities that can be represented mathematically by vectors.



## Vector Denoting

- Vectors are writing with an arrow on top on equations.


Ex


- Velocity vector  $\vec{v}$

- Force vector  $\vec{F}$

- ⊗ Note Any variable symbol with no arrow on top means scalar.

- A vector can be geometrically represented by a direction line segment with a head & a tail;



- So vector  $\vec{AB}$  is a vector from point A to B.
- Also, we can denote vector  $\vec{AB}$  by a small case letter  $\vec{v}$
- The length of the arrow  corresponds to the magnitude of the vector.
- The arrow points in the direction of the vector.




How to represent the vector mathematically

Vector in plane :-  $\vec{v}$  (متجه في المستوى)

We can write vectors as Columns. Let us take a very important special vector as example :-

$$i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad j = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$




Any vector in the xy-plane can be written in terms of  $i$  &  $j$  using the triangle law & scalar multiplication.

$$\vec{v} = ai + bj = a \begin{pmatrix} 1 \\ 0 \end{pmatrix} + b \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$$


Ex1

$$\begin{pmatrix} 3 \\ 2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 2 \begin{pmatrix} 0 \\ 1 \end{pmatrix} = 3i + 2j$$



Notes

\* If  $a=b=1$ , then  $\vec{v} = i + j$  is a "unit vector"






Finding the length/magnitude and the direction of vector  $\vec{v}$  إيجاد طول (مقدار) واتجاه المتجه

If  $\vec{v} = a\mathbf{i} + b\mathbf{j}$  (1), then the length/magnitude of a vector  $\vec{v}$  is -

$|\vec{v}| = \sqrt{a^2 + b^2}$  (2)  
 (نظرية فيثاغورس)

• It's a Pythagorean theorem



$a = |\vec{v}| \cos \alpha$   
 $b = |\vec{v}| \sin \alpha$  (3)

$\tan \alpha = \frac{b}{a}$

Substitute eq. (3) in (1) yields ;

$\vec{v} = |\vec{v}| (\cos \alpha \mathbf{i} + \sin \alpha \mathbf{j})$

- $\vec{v}$  ~ vector symbol
- $|\vec{v}|$  ~ vector length
- $\mathbf{i}, \mathbf{j}$  ~ unit vector components (basis / Fundamental vector components)
- $\alpha$  ~ vector angle with x-axis

Ex) Find a vector in plane of length (7 units) & makes angle (35°) with x-axis?

Solution

since  $|\vec{v}| = 7$  &  $\alpha = 35^\circ$

$\therefore \vec{v} = 7 * (\cos 35 \mathbf{i} + \sin 35 \mathbf{j})$

$\vec{v} = 5.7 \mathbf{i} + 4 \mathbf{j}$  Ans




Ex) Find the angle between the vector  
 $\vec{v} = 2\hat{i} + 3\hat{j}$  and the x-axis?

Solution

$$|\vec{v}| = \sqrt{a^2 + b^2} \quad ; \quad a = 2$$

$$b = 3$$

$$|\vec{v}| = \sqrt{2^2 + 3^2} = \sqrt{13}$$



$$a = |\vec{v}| \cos \alpha \Rightarrow \cos \alpha = \frac{a}{|\vec{v}|} = \frac{2}{\sqrt{13}}$$


$$\therefore \alpha = \cos^{-1} \frac{2}{\sqrt{13}} = \boxed{56.3^\circ} \quad \underline{\text{Ans}}$$

Vectors with tail not in origin so  
(a,0)  $\hat{i}$  and (0,b)  $\hat{j}$   $\in$   $\hat{i}$   $\hat{j}$   $\in$   $\hat{i}$   $\hat{j}$

Vectors can be start not from the origin, but from any where like A to B

$$\therefore \vec{AB} = \Delta a \hat{i} + \Delta b \hat{j}$$

$$\vec{AB} = (a_2 - a_1) \hat{i} + (b_2 - b_1) \hat{j}$$






## Vectors in a space

This could be in three or higher dimensions

Similar to the 2-D case,  
 but we now have three  
 basis vectors  $i, j$  &  $k$ .

From these three  
 components unit vectors  
 we can describe any  
 vector in three-dimensional  
 space.



$$\vec{v} = \vec{OA} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$$

where  $\mathbf{i}$

$\mathbf{i}, \mathbf{j}, \mathbf{k}$  = Basis or Fundamental unit vector.

$a, b, c$  = Directional numbers (scalars).

$\alpha, \beta, \gamma$  = Directional angles.

$$|\vec{v}| = |\vec{OA}| = \sqrt{a^2 + b^2 + c^2}$$

$$a = |\vec{v}| \times \cos \alpha$$

$$b = |\vec{v}| \times \cos \beta$$

$$c = |\vec{v}| \times \cos \gamma$$

$$\frac{\vec{v}}{|\vec{v}|} = \cos \alpha \mathbf{i} + \cos \beta \mathbf{j} + \cos \gamma \mathbf{k} \quad \left. \vphantom{\frac{\vec{v}}{|\vec{v}|}} \right\} = \text{unit vector in the direction of } \vec{v}$$

And,

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$



Ex1 Find a vector in space of length (5 units) that makes angles ( $70^\circ$ ) with x-axis, ( $85^\circ$ ) with y-axis ?

Solution

$$\alpha = 70^\circ, \beta = 85^\circ, |\vec{v}| = 5$$
$$\gamma = ?, \vec{v} = ?$$

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1 \Rightarrow \cos^2 70^\circ + \cos^2 85^\circ + \cos^2 \gamma = 1$$

$$\therefore \boxed{\cos \gamma = 0.935}$$

$$\vec{v} = |\vec{v}| (\cos \alpha \hat{i} + \cos \beta \hat{j} + \cos \gamma \hat{k})$$
$$= 5 (\cos 70^\circ \hat{i} + \cos 85^\circ \hat{j} + 0.935 \hat{k})$$

$$\boxed{\vec{v} = 1.7 \hat{i} + 0.436 \hat{j} + 4.675 \hat{k}}$$

Ans

Ex2 Find the angle between the vector  $\vec{v} = -4\hat{i} + 5\hat{j} + \hat{k}$  and the x-axis?

Solution


$$a = -4, b = 5, c = 1$$

$$|\vec{v}| = \sqrt{a^2 + b^2 + c^2}$$

$$|\vec{v}| = \sqrt{(-4)^2 + (5)^2 + (1)^2} = \sqrt{42}$$

$$\cos \alpha = \frac{a}{|\vec{v}|} \Rightarrow \alpha = \cos^{-1} \frac{a}{|\vec{v}|} = \cos^{-1} \frac{-4}{\sqrt{42}}$$

$$\boxed{\alpha = 128^\circ}$$





## Scalar product = (Dot Product) ضرب النقط المبرية

Let  $\vec{A} = a_1\vec{i} + a_2\vec{j} + a_3\vec{k}$

And  $\vec{B} = b_1\vec{i} + b_2\vec{j} + b_3\vec{k}$

Then,  $\vec{A} \cdot \vec{B} = |\vec{A}| |\vec{B}| \cos \theta$

Where  $\theta$  is the angle between  $\vec{A}$  &  $\vec{B}$

### Properties

1-  $\vec{A} \cdot \vec{A} = |\vec{A}|^2$

2-  $\vec{A} \cdot \vec{B} = \vec{B} \cdot \vec{A} = a_1b_1 + a_2b_2 + a_3b_3$

3-  $\cos \theta = \frac{\vec{A} \cdot \vec{B}}{|\vec{A}| |\vec{B}|} = \frac{a_1b_1 + a_2b_2 + a_3b_3}{\sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}}$

4-  $\vec{A} \perp \vec{B} \Rightarrow \vec{A} \cdot \vec{B} = 0$  [Orthogonal Vectors]  
متعامدتين

5-  $a_1\vec{i} + b_2\vec{j} \perp b_1\vec{i} - a_2\vec{j}$

EX) Find the angle  $\theta$  between  $\vec{A} = \vec{i} - 2\vec{j} - 2\vec{k}$  &  $\vec{B} = 6\vec{i} + 3\vec{j} + 2\vec{k}$ ?

Solution

$\vec{A} \cdot \vec{B} = (1 \times 6) + (-2 \times 3) + (-2 \times 2) = \boxed{-4}$

$|\vec{A}| = \sqrt{1^2 + (-2)^2 + (-2)^2} = \sqrt{9} = \boxed{3}$

$|\vec{B}| = \sqrt{6^2 + 3^2 + 2^2} = \sqrt{49} = \boxed{7}$

$|\vec{A}| |\vec{B}| = \boxed{21}$

$\cos^{-1} \frac{\vec{A} \cdot \vec{B}}{|\vec{A}| |\vec{B}|} = \cos^{-1} \frac{-4}{21} \approx \boxed{101^\circ}$  Ans






## Vector Product :- (Cross-product) حرب المتجهات

Normal vector is what yields from vector product or cross product -

$$\vec{N} = \vec{A} \times \vec{B} = \vec{n} |\vec{A}| |\vec{B}| \sin \theta$$

where  $\vec{n}$  is a normal unit vector



$$\vec{A} \times \vec{B} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} ; \text{ where, } \vec{A} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k} \\ \vec{B} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$$

### Properties :-

- $\vec{A} \times \vec{A} = 0 \rightarrow \text{"sin } 0 = 0\text{"}$
- $\vec{A} \times \vec{B} = -\vec{B} \times \vec{A}$
- $\vec{A} \parallel \vec{B} \Rightarrow \vec{A} \times \vec{B} = 0 \Rightarrow \text{sin } 0 = 0$
- Area of  $\Delta ABC = \frac{1}{2} |\vec{A} \times \vec{B}|$



Ex) Find  $\vec{A} \times \vec{B}$  &  $\vec{B} \times \vec{A}$  if

$$\vec{A} = 2\vec{i} + \vec{j} + \vec{k}$$

$$\vec{B} = -4\vec{i} + 3\vec{j} + \vec{k}$$

Solutions

$$\vec{A} \times \vec{B} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & 1 & 1 \\ -4 & 3 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 1 \\ 3 & 1 \end{vmatrix} \vec{i} - \begin{vmatrix} 2 & 1 \\ -4 & 1 \end{vmatrix} \vec{j} + \begin{vmatrix} 2 & 1 \\ -4 & 3 \end{vmatrix} \vec{k}$$

$$= (1 \times 1 - (3 \times 1))\vec{i} - (2 \times 1 - (-4 \times 1))\vec{j} + (2 \times 3 - (-4 \times 1))\vec{k}$$

$$\boxed{\vec{A} \times \vec{B} = -2\vec{i} - 6\vec{j} + 10\vec{k}}$$

but,

$$\vec{B} \times \vec{A} = -\vec{A} \times \vec{B} = \boxed{2\vec{i} + 6\vec{j} - 10\vec{k}}$$

Triple Product :- الضرب الثلاثي الكروي

A- Scalar triple product :-

$$\boxed{\vec{A} \cdot (\vec{B} \times \vec{C}) = (\vec{A} \times \vec{B}) \cdot \vec{C}}$$

Note

1- Box volume is  $\Rightarrow V_{\text{box}} = |\vec{A} \cdot \vec{B} \times \vec{C}|$

2- Pyramid volume is  $\Rightarrow V_{\text{py}} = \frac{1}{6} |\vec{A} \cdot \vec{B} \times \vec{C}|$

B- Vector triple product :-

$$\boxed{\vec{A} \times (\vec{B} \times \vec{C}) = (\vec{A} \cdot \vec{C})\vec{B} - (\vec{A} \cdot \vec{B})\vec{C}}$$

Note

CS Scanned with CamScanner;  $\vec{i} \cdot \vec{j} = \vec{j} \cdot \vec{k} = \vec{k} \cdot \vec{i} = 1$ ;  $\vec{i} \times \vec{i} = \vec{j} \times \vec{j} = \vec{k} \times \vec{k} = 0$




$$\hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0$$

$$\hat{i} \times \hat{j} = \hat{k}$$

$$\hat{j} \times \hat{k} = \hat{i}$$

$$\hat{k} \times \hat{i} = \hat{j}$$



## HW#2

1- Find the length & direction of these vectors & the angles make with the x-axis?

$$a - 5\hat{i} + 12\hat{j}$$

$$b - \sqrt{3}\hat{i} + \hat{j}$$

2- Find a vector 6 units long in the direction of  $\vec{A} = 2\hat{i} + 2\hat{j} - \hat{k}$

3- Find the area of the triangle whose vertices are  $A(1, -1, 0)$ ,  $B(2, 1, -1)$ ,  $C(-1, 1, 2)$ ?

4- If  $\vec{A} = 2\hat{i} - \hat{j}$  &  $\vec{B} = \hat{i} + 3\hat{j} - 2\hat{k}$ ,  
Find  $\vec{A} \times \vec{B}$ , then calculate  $(\vec{A} \times \vec{B}) \cdot \vec{A}$ ?

---- نهاية محاضرة " Vectors, Vectors in Space, Unit Vector, Scalar Product, Vector Product المتجهات، المتجهات في الفضاء، وحدة المتجه، ضرب القيمة العددية، ضرب المتجه " ----