



Class: 2<sup>st</sup>

Subject: Mathematics

Lecturer: M.Sc Murtadha Mohsen Al-Masoudy

E-mail: [Murtadha\\_Almasoody@mustaqbal-college.edu.iq](mailto:Murtadha_Almasoody@mustaqbal-college.edu.iq)



M.Sc. Murtadha Al-Masoudy  
Mathematics-2-  
2<sup>nd</sup> Year

### Multiple Integrals

The multiple integrals are the integrals of a function of two or more variables over a region in the plane or space.

#### Double Integrals :-

① If a function  $f(x,y)$  is defined on a rectangular region  $R$

$R: a \leq x \leq b \ \& \ c \leq y \leq d$ , then

$$\iint_R f(x,y) dA = \int_c^d \int_a^b f(x,y) dx dy = \int_a^b \int_c^d f(x,y) dy dx$$

② If a function  $f(x,y)$  is defined on a region  $R$

$R: a \leq x \leq b \ \& \ f_1(x) \leq y \leq f_2(x)$ , then

$$\iint_R f(x,y) dA = \int_a^b \int_{f_1(x)}^{f_2(x)} f(x,y) dy dx$$

③ If a function  $f(x,y)$  is defined on a region  $R$

$R: g_1(x) \leq x \leq g_2(x) \ \& \ c \leq y \leq d$

$$\iint_R f(x,y) dA = \int_c^d \int_{g_1(x)}^{g_2(x)} f(x,y) dx dy$$



Ex: Calculate  $\iint_R f(x,y) dA$  for

$$f(x,y) = 1 - 6x^2y \quad \& \quad R: 0 \leq x \leq 2, -1 \leq y \leq 1$$

Solution: —

$$\begin{aligned} \iint_R f(x,y) dA &= \int_0^2 \int_{-1}^1 (1 - 6x^2y) dy dx = \int_0^2 \left[ y - \frac{6x^2y^2}{2} \right]_{-1}^1 dx \\ &= \int_0^2 \left[ \left(1 - \frac{6x^2 \cdot 1^2}{2}\right) - \left(-1 - \frac{6x^2(-1)^2}{2}\right) \right] dx \\ &= \int_0^2 [1 - 3x^2 - (-1 - 3x^2)] dx = \int_0^2 2 dx = 2x \Big|_0^2 \\ &= (2 \cdot 2) - (2 \cdot 0) = \boxed{4} \end{aligned}$$

Ex: Calculate  $\iint_0^1 e^{y^2} dx \cdot dy$ .

Solution: —

$$\begin{aligned} \iint_0^1 e^{y^2} dx \cdot dy &= \int_0^1 [e^{y^2} x]_0^1 dy = \int_0^1 [(e^{y^2} \cdot 1) - (e^{y^2} \cdot 0)] dy \\ &= \int_0^1 y e^{y^2} dy \cdot \frac{2}{2} = \frac{1}{2} [e^{y^2}]_0^1 = \frac{1}{2} (e^1 - e^0) \\ &= \boxed{\frac{1}{2} (e^1 - 1)} \end{aligned}$$

Ex: Find  $\int_0^2 \int_{y^2}^{6-y} dx \cdot dy$

Solution: —

$$\begin{aligned} \int_0^2 \int_{y^2}^{6-y} dx \cdot dy &= \int_0^2 [x]_{y^2}^{6-y} dy = \int_0^2 (6 - y - y^2) dy = 6y - \frac{y^2}{2} - \frac{y^3}{3} \Big|_0^2 \\ &= \left(6 \cdot 2 - \frac{2^2}{2} - \frac{2^3}{3}\right) - (0) = 12 - 2 - \frac{8}{3} \\ &= \frac{22}{3} = \boxed{7.33} \end{aligned}$$





Class: 2<sup>st</sup>

Subject: Mathematics

Lecturer: M.Sc Murtadha Mohsen Al-Masoudy

E-mail: [Murtadha\\_Almasoody@mustaqbal-college.edu.iq](mailto:Murtadha_Almasoody@mustaqbal-college.edu.iq)



Ex: - Calculate  $\iint_R f(x,y) dA$  for

$$f(x,y) = 4x + 2 \quad \& \quad R: x^2 \leq y \leq 2x, \quad 0 \leq x \leq 2$$

Solution: -

$$\iint_R f(x,y) dA = \int_0^2 \int_{x^2}^{2x} (4x+2) dy dx = \int_0^2 (4xy + 2y) \Big|_{y=x^2}^{y=2x} dx$$

$$= \int_0^2 [(4x \cdot 2x + 2 \cdot 2x) - (4x \cdot x^2 + 2 \cdot x^2)] dx$$

$$= \int_0^2 [8x^2 + 4x - (4x^3 + 2x^2)] dx = \int_0^2 (6x^2 + 4x - 4x^3) dx$$

$$= \left[ \frac{6x^3}{3} + \frac{4x^2}{2} - \frac{4x^4}{4} \right]_0^2 = \left[ 2x^3 + 2x^2 - x^4 \right]_0^2$$

$$= 2(2)^3 + 2(2)^2 - (2)^4 = 16 + 8 - 16 = \boxed{8}$$

Ex: - Evaluate the  $\iint_R \frac{\sin x}{x} dA$  where  $R$  is the triangle in  $xy$ -plane bounded by the  $x$ -axis, the line  $x=y$  and the line  $x=1$ .

Solution: -

$$\int_0^1 \int_0^x \frac{\sin x}{x} dy dx = \int_0^1 \frac{\sin x}{x} \Big|_{y=0}^{y=x} dx$$

$$\int_0^1 \frac{\sin x}{x} \cdot x dx = -\cos x \Big|_0^1$$

$$= -(\cos(1) - \cos(0))$$

$$= \boxed{0.459}$$

