

Solubility of Weak Electrolytes

Many important drugs belong to the class of weak acids and bases.

Weak acids react with dilute alkalis to form water-soluble salts, but they can be precipitated as the free acids if stronger acidic substances are added to the solution. For example, a 1% solution of phenobarbital sodium is soluble at pH values high in the alkaline range. The soluble ionic form is converted into molecular phenobarbital as the pH is lowered, and below 9.3, the drug begins to precipitate from solution.

Weak bases react with dilute acids (decrease the pH) to form water-soluble salts, but they can be precipitated as the free bases if stronger basic substances (increase the pH) are added to the solution.

For example, alkaloid salts such as atropine sulfate begin to precipitate as the pH is elevated.

To ensure a clear homogeneous solution and maximum therapeutic effectiveness, the preparations should be adjusted to an optimum pH.

Calculating the Solubility of Weak Electrolytes as Influenced by pH

According to the Henderson-Hasselbach equation, the relationship between pH, pKa, and relative concentrations of an acid and its salt is as follows:

$$\text{pH} = \text{pKa} + \log \frac{[A^-]}{[HA]}$$

where $[A^-]$ is the molar concentration of the salt (dissociated species) and $[HA]$ is the concentration of the undissociated acid. When the concentrations of salt and acid are equal, the pH of the system equals the pKa of the acid.

As the pH decreases, the concentration of the molecular acid increases and that of the salt decreases.

Changes in solubility brought about by alterations of solvent pH can be predicted by the pHp equation. The pHp is the pH below which an acid or above which a base will begin to precipitate.

$$\text{pH} = \text{pK}_a + \log \frac{S - S_0}{S_0} \quad (\text{For weak acid})$$

$$\text{pH} = \text{pK}_w - \text{pK}_b + \log \frac{S_0}{S - S_0} \quad (\text{For weak base})$$

where,

- S_0 = the molar solubility of the undissociated acid or base
- S = the molar concentration of the salt form of the drug initially added

Example 10.16

Below what pH will free phenobarbital begin to separate from a solution having an initial concentration of 1 g of sodium phenobarbital per 100 mL at 25°C? The molar solubility, S_0 , of phenobarbital is 0.0050 and the pK_a is 7.41 at 25°C. The molecular weight of sodium phenobarbital is 254.

The molar concentration of salt initially added is
(g/liter)/m.wt= 10/254= 0.039mole/liter

$$\text{pH}_p = 7.41 + \log \frac{(0.039 - 0.005)}{0.005} = 8.24$$

Q38. The molar solubility of sulfathiazole (weak acid) in water is 0.002, the $pK_a=7.12$, m.wt of sodium sulfathiazole = 304, what is the lowest pH allowable for complete solubility in a 5% solution of salt?

$$pK_a = 7.12 \quad S_0 = 0.002$$

$$M \text{ salt} = \frac{wt}{m.wt} \times \frac{1000}{vol}$$

$$= \frac{5}{304} \times \frac{1000}{100} = 0.164$$

$$pH_p = 7.12 + \log \frac{(0.164 - 0.002)}{0.002} = 9.03$$

The Influence of Solvents on the Solubility of Drugs

Weak electrolytes can behave like strong electrolytes or like nonelectrolytes in solution.

When the solution is of such a pH that the drug is entirely in the ionic form, it behaves as a solution of a strong electrolyte-----> no problem.

However, when the pH is adjusted to a value at which un-ionized molecules are produced in sufficient concentration to exceed the solubility of this form, precipitation occurs. -----> (Problem)

To solve this problem, a solute is more soluble in a mixture of solvents than in one solvent alone. This phenomenon is known as cosolvency, and the solvents that, in combination, increase the solubility of the solute are called cosolvents.

For example phenobarbital solubility is increased when add alcohol or glycerin to water.

Combined Effect of pH and Solvents

The solvent affects the solubility of a weak electrolyte in a buffered solution in two ways:

(a) The addition of alcohol to a buffered aqueous solution of a weak electrolyte increases the solubility of the un-ionized species by adjusting the polarity of the solvent to a more favorable value.

(b) Because it is less polar than water, alcohol decreases the dissociation of a weak electrolyte, and the solubility of the drug goes down as the dissociation constant is decreased (pKa is increased).

Influence of Surfactants

Weakly acidic and basic drugs can be brought into solution by the solubilizing action of surface-active agents such as detergent.

Influence of complexation

Complexation may be used to increase the solubility such as addition of potassium iodide to iodine.

Sometimes complexation causes a decrease in solubility such as the complex between tetracycline and calcium produces an insoluble complex.

Influence of size and shape of particles


The size of particles affects solubility. Solubility increases with decreasing particle size as a consequence of an increase in surface area.

The configuration of a molecule and the type of arrangement in the crystal also has some influence on solubility, and a symmetric particle can be less soluble than an unsymmetrical one.

Solubility of Slightly Soluble Electrolytes

When slightly soluble electrolytes are dissolved to form saturated solutions, the solubility is described by a special constant, known as the solubility product, K_{sp} , of the compound.

Silver chloride is an example of such a slightly soluble salt. The excess solid in equilibrium with the ions in saturated solution at a specific temperature is represented by the equation




and because the salt dissolves only with difficulty and the ionic strength is low, the equilibrium expression can be written in terms of concentrations instead of activities:

$$\frac{[\text{Ag}^+][\text{Cl}^-]}{[\text{AgCl}_{\text{solid}}]} = K$$

Moreover, because the concentration of the solid phase is essentially constant,

$$[\text{Ag}^+][\text{Cl}^-] = K_{sp}$$

As in the case of other equilibrium expressions, the concentration of each ion is raised to a power equal to the number of ions appearing in the formula. Thus, for aluminum hydroxide, $\text{Al}(\text{OH})_3$.





EXAMPLE 10-13: The measured solubility of silver chloride in water at 20°C is 1.12×10^{-5} mole/liter. This is also the concentration of the silver ion and the chloride ion because silver chloride is nearly completely dissociated. Calculate the solubility product of this salt. We have

$$K_{sp} = (1.12 \times 10^{-5}) \times (1.12 \times 10^{-5}) = 1.25 \times 10^{-10}$$

If an ion in common with AgCl, that is, Ag^+ or Cl^- , is added to a solution of silver chloride, the equilibrium is altered.

The addition of sodium chloride, for example, increases the concentration of chloride ions so that momentarily $[Ag^+][Cl^-] > K_{sp}$ and some of the AgCl precipitates from the solution (the reaction shift to left) until the equilibrium $[Ag^+][Cl^-] = K_{sp}$ is reestablished. Hence, the result of adding a common ion is to reduce the solubility of a slightly soluble electrolyte

Salts having no ion in common with the slightly soluble electrolyte produce an effect opposite to that of a common ion: At moderate concentration, they increase rather than decrease the solubility because they lower the activity coefficient.

Distribution of Solutes between Immiscible Solvents

If an excess of substance is added to a mixture of two immiscible liquids, it will distribute itself between the two phases so that each becomes saturated.

If the substance is added to the immiscible solvents in an amount insufficient to saturate the solutions, it will still become distributed between the two layers in a definite concentration ratio.

If C_1 and C_2 are the equilibrium concentrations of the substance in Solvent1 and Solvent2, respectively, the equilibrium expression becomes

$$K = \frac{C_1}{C_2}$$

The above equation is known as the **distribution law**

The equilibrium constant, K , is known as the distribution ratio, distribution coefficient, or partition coefficient.

The partition law states that:

- At a given temperature, the ratio of the concentrations of a solute in two immiscible solvents (solvent 1 and solvent 2) is constant when equilibrium has been reached
- This constant is known as the partition coefficient (or distribution coefficient)

Example 9-5

When boric acid is distributed between water and amyl alcohol at 25°C, the concentration in water is found to be 0.0510 mole/liter and in amyl alcohol it is found to be 0.0155 mole/liter. What is the distribution coefficient? We have

$$K = \frac{C.H_2O}{C.alc} = \frac{0.051}{0.0155} = 3.29$$

No convention has been established with regard to whether the concentration in the water phase or that in the organic phase should be placed in the numerator.

Therefore, the result can also be expressed as

$$K = \frac{C.alc}{C.H_2O} = \frac{0.0155}{0.0510} = 0.304$$

One should always specify, which of these two ways the distribution constant is being expressed.

Importance of partition Knowledge

The principle is involved in several areas:

- 1- Drugs partitioning between aqueous phases and lipid biophases
- 2- Preservation of oil–water systems
- 3- Absorption and distribution of drugs throughout the body
- 4- Antibiotics partitioning into microorganisms
- 5- Solvent extraction
- 6- Chromatography


Partition law holds true

- 1- at constant temperature
- 2- when the solute exists in the same form in both solvents (the species are common to both phases)


Effect of Ionic Dissociation and Molecular Association on Partition

Some compounds have more than one species in oil phase and in aqueous phase which produce complicated case in calculation of partition coefficient.

Example, benzoic acid that is used as preservative present in oil phase as monomer and as dimer in equilibrium.



Benzoic acid present in aqueous phase as unionized (HA) and as ionized (A^-) in equilibrium.



The true distribution coefficient, K is the ratio of molar concentration of the species common to both the oil and water phases

$$K = \frac{[\text{HA}]_o}{[\text{HA}]_w}$$

The experimentally observed or apparent distribution coefficient determined by using the total acid concentration (all species) obtained by analysis is

$$K' = \frac{[\text{HA}]_o + [\text{HA}]_d}{[\text{HA}]_w + [\text{A}^-]_w}$$

Thus the observed distribution coefficient depends on two equilibria.

In the oil phase, the concentration of monomer or dimer depends on the type of oil, for example benzoic only present as monomer in peanut oil.

In the aqueous phase, the concentration of unionized or ionized depends on the pKa of compound and pH of solvent.

