


2nd region

$$\left. \begin{array}{l} y = 1 - x \dots(1) \\ y = -\frac{x}{2} \dots(2) \end{array} \right\} \Rightarrow x = 2 \Rightarrow y = -1 \quad y \text{ from } 0 \text{ to } 2$$

$$\begin{aligned} \int_0^2 \int_{-\frac{y}{2}}^{1-y} dx dy + \int_{-1}^0 \int_{-2y}^{1-y} dx dy &= \int_0^2 x \Big|_{-\frac{y}{2}}^{1-y} dy + \int_{-1}^0 x \Big|_{-2y}^{1-y} dy \\ &= \int_0^2 \left(1 - \frac{y}{2}\right) dy + \int_{-1}^0 (1 + y) dy = y - \frac{y^2}{4} \Big|_0^2 + y + \frac{y^2}{2} \Big|_{-1}^0 \\ &= 2 - 1 - 0 + 0 - \left(-1 + \frac{1}{2}\right) = \frac{3}{2} \\ &= \text{The same result as in (a).} \end{aligned}$$




### 7-3- Triple integrals (Volume):

Consider a region  $N$  in  $xyz$ -space bounded below by a surface  $z = f_1(x, y)$ , above by the surface  $z = f_2(x, y)$  and laterally by a cylinder  $c$  with elements parallel to the  $z$ -axis. Let  $A$  denote the region of the  $xy$ -plane enclosed by cylinder  $c$  (that is,  $A$  is the region covered by the orthogonal projection of the solid into  $xy$ -plane). Then the volume  $V$  of the region  $V$  can be found by evaluating the triply iterated integral:-

$$V = \iint_A \int_{f_1(x,y)}^{f_2(x,y)} dz dy dx$$

Let  $z$ -limits of integration indicate that for every  $(x, y)$  in the region  $A$ ,  $z$  may extend from the lower surface  $z = f_1(x, y)$  to the surface  $z = f_2(x, y)$ . The  $y$ - and  $x$ -limits of integration have not been given explicitly in equation above, but are indicated as extending over the region  $A$ .



We can find the equation of the boundary of the region  $A$  by eliminating  $z$  between the two equations  $z = f_1(x, y)$  and  $z = f_2(x, y)$ , thus obtaining an equation  $f_1(x, y) = f_2(x, y)$  which contains no  $z$ , and interpret it as an equation in the  $xy$ -plane.

**EX-9** The volume in the first octant bounded by the cylinder  $x = 4 - y^2$ , and the planes  $z = y$ ,  $x = 0$ ,  $z = 0$ .

**Sol.-**

$$x = 4 - y^2 \Rightarrow y = \pm\sqrt{4-x} \quad \text{in first octant :-}$$

$$\begin{aligned} V &= \int_0^4 \int_0^{\sqrt{4-x}} \int_0^y dz \, dy \, dx = \int_0^4 \int_0^{\sqrt{4-x}} z \Big|_0^y \, dy \, dx = \int_0^4 \int_0^{\sqrt{4-x}} y \, dy \, dx = \int_0^4 \frac{y^2}{2} \Big|_0^{\sqrt{4-x}} \, dx \\ &= \frac{1}{2} \int_0^4 (4-x-0) \, dx = \frac{1}{2} \left[ 4x - \frac{x^2}{2} \right]_0^4 = \frac{1}{2} \left[ 16 - \frac{16}{2} - 0 \right] = 4 \end{aligned}$$

**EX-10** The volume enclosed by the cylinders  $z = 5 - x^2$ ,  $z = 4x^2$  and the planes  $y = 0$ ,  $x + y = 1$ .

**Sol.-**

$$\left. \begin{aligned} z &= 5 - x^2 \dots(1) \\ z &= 4x^2 \dots\dots(2) \end{aligned} \right\} \Rightarrow x = \pm 1$$

$$\begin{aligned} V &= \int_{-1}^1 \int_0^{1-x} \int_{4x^2}^{5-x^2} dz \, dy \, dx = \int_{-1}^1 \int_0^{1-x} z \Big|_{4x^2}^{5-x^2} \, dy \, dx = \int_{-1}^1 \int_0^{1-x} (5 - 5x^2) \, dy \, dx \\ &= 5 \int_{-1}^1 (1-x^2) y \Big|_0^{1-x} \, dx = 5 \int_{-1}^1 (1-x^2)(1-x) \, dx \\ &= 5 \int_{-1}^1 (1-x-x^2+x^3) \, dx = 5 \left[ x - \frac{x^2}{2} - \frac{x^3}{3} + \frac{x^4}{4} \right]_{-1}^1 \\ &= 5 \left[ (1+1) - \frac{1}{2}(1-1) - \frac{1}{3}(1+1) + \frac{1}{4}(1-1) \right] = \frac{20}{3} \end{aligned}$$

**EX-11** The volume enclosed by the cylinders  $y^2 + 4z^2 = 16$  and the planes  $x = 0$ ,  $x + y = 4$ .

**Sol.-**

$$y^2 + 4z^2 = 16 \Rightarrow y = \pm 2\sqrt{4 - z^2}$$

$$\begin{aligned} V &= \int_{-2}^2 \int_{-2\sqrt{4-z^2}}^{2\sqrt{4-z^2}} \int_0^{4-y} dx dy dz \\ &= \int_{-2}^2 \int_{-2\sqrt{4-z^2}}^{2\sqrt{4-z^2}} (4 - y) dy dz = \int_{-2}^2 4y - \frac{y^2}{2} \Big|_{-2\sqrt{4-z^2}}^{2\sqrt{4-z^2}} dz = 16 \int_{-2}^2 (4 - z^2)^{1/2} dz \end{aligned}$$

$$\text{let } z = 2 \sin \theta \Rightarrow dz = 2 \cos \theta d\theta, \quad \theta = \sin^{-1} \frac{z}{2} \quad \begin{matrix} \text{at } z=2 \Rightarrow \theta = \frac{\pi}{2} \\ \Rightarrow \Rightarrow \Rightarrow \\ \text{at } z=-2 \Rightarrow \theta = \frac{\pi}{2} \end{matrix}$$

$$V = 16 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (4 - 4 \sin^2 \theta)^{1/2} 2 \cos \theta d\theta = 64 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^2 \theta d\theta = 64 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{1 + \cos 2\theta}{2} d\theta$$

$$= 32 \left[ \theta + \frac{1}{2} \sin 2\theta \right]_{-\frac{\pi}{2}}^{\frac{\pi}{2}} = 32 \left[ \left( \frac{\pi}{2} + \frac{\pi}{2} \right) + \frac{1}{2} (0 - 0) \right] = 32\pi$$

**EX-12** The volume bounded by the ellipse paraboloids  $z = x^2 + 9y^2$  and  $z = 18 - x^2 - 9y^2$ .

**Sol.-**

$$\left. \begin{aligned} z &= 18 - x^2 - 9y^2 \quad \dots(1) \\ z &= x^2 + 9y^2 \quad \dots\dots\dots(2) \end{aligned} \right\} \Rightarrow 9 - x^2 - 9y^2 = 0 \Rightarrow y = \pm \frac{1}{3} \sqrt{9 - x^2}$$

$$V = \int_{-3}^3 \int_{-\frac{1}{3}\sqrt{9-x^2}}^{\frac{1}{3}\sqrt{9-x^2}} \int_{x^2+9y^2}^{18-x^2-9y^2} dz dy dx = \int_{-3}^3 \int_{-\frac{1}{3}\sqrt{9-x^2}}^{\frac{1}{3}\sqrt{9-x^2}} [18 - x^2 - 9y^2 - (x^2 + 9y^2)] dy dx$$

$$\begin{aligned}
 V &= 2 \int_{-3}^3 (9-x^2)y - 3y^3 \Bigg|_{-\frac{1}{3}\sqrt{9-x^2}}^{\frac{1}{3}\sqrt{9-x^2}} dx \\
 &= 2 \int_{-3}^3 \left[ (9-x^2) \left( \frac{\sqrt{9-x^2}}{3} + \frac{\sqrt{9-x^2}}{3} \right) - 3 \left( \frac{(9-x^2)^{3/2}}{27} + \frac{(9-x^2)^{3/2}}{27} \right) \right] dx \\
 &= \frac{8}{9} \int_{-3}^3 (9-x^2)^{3/2} dx
 \end{aligned}$$

let  $x = 3\sin\theta \Rightarrow dx = 3\cos\theta d\theta$  ,  $\theta = \sin^{-1} \frac{x}{3}$    
at  $x=3 \Rightarrow \theta = \frac{\pi}{2}$   
 $\Rightarrow \Rightarrow \Rightarrow$   
at  $x=-3 \Rightarrow \theta = -\frac{\pi}{2}$

$$\begin{aligned}
 &= \frac{8}{9} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (9-9\sin^2\theta)^{3/2} 3\cos\theta d\theta = 72 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^4\theta d\theta = 72 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \left( \frac{1+\cos 2\theta}{2} \right)^2 d\theta \\
 &= 18 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (1+2\cos 2\theta + \cos^2 2\theta) d\theta = 18 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \left( 1+2\cos 2\theta + \frac{\cos 4\theta}{2} \right) d\theta \\
 &= 9 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (3+4\cos 2\theta + \cos 4\theta) d\theta = 9 \left[ 3\theta + 2\sin 2\theta + \frac{1}{4}\sin 4\theta \right]_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \\
 &= 9 \left[ 3\left(\frac{\pi}{2} + \frac{\pi}{2}\right) + 2(\sin \pi - \sin(-\pi)) + \frac{1}{4}(\sin 2\pi - \sin(-2\pi)) \right] = 27\pi
 \end{aligned}$$


#### 7-4- The length of a plane curve:-

The length of the curve  $y = f(x)$  from point  $A(a,c)$  to  $B(b,d)$  is:-

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

If  $x$  can be expressed as a function of  $y$  then the length is:-

$$L = \int_c^d \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$



Let the equation of motion be  $x = g(t)$  and  $y = h(t)$  continuously differentiable for  $t$  between  $t_a$  (at A) and  $t_b$  (at B), then the length of the curve is:-

$$L = \int_{t_a}^{t_b} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

**EX-13** – Find the length of the curve:

1)  $y = \frac{1}{3}(x^2 + 2)^{\frac{3}{2}}$  from  $x = 0$  to  $x = 3$

2)  $9x^2 = 4y^3$  from  $(0,0)$  to  $(2\sqrt{3},3)$

3)  $y = x^{\frac{2}{3}}$  from  $x = -1$  to  $x = 8$

**Sol.**

1)  $y = \frac{1}{3}(x^2 + 2)^{\frac{3}{2}} \Rightarrow \frac{dy}{dx} = x(x^2 + 2)^{\frac{1}{2}}$

$$L = \int_0^3 \sqrt{1 + x^2(x^2 + 2)} dx = \int_0^3 (x^2 + 1) dx = \left. \frac{x^3}{3} + x \right|_0^3 = 9 + 3 - 0 = 12$$

2)  $9x^2 = 4y^3 \Rightarrow x = \mp \frac{2}{3}y^{\frac{3}{2}}$  Since  $x$  from 0 to  $2\sqrt{3}$


then  $x = \frac{2}{3}y^{\frac{3}{2}} \Rightarrow \frac{dx}{dy} = y^{\frac{1}{2}}$

$$L = \int_0^3 \sqrt{1 + y} dy = \left. \frac{2}{3}(1 + y)^{\frac{3}{2}} \right|_0^3 = \frac{2}{3}[8 - 1] = \frac{14}{3}$$

3)  $y = x^{\frac{2}{3}} \Rightarrow \frac{dy}{dx} = \frac{2}{3}x^{-\frac{1}{3}}$

Since  $\frac{dy}{dx} = \infty$  at  $x = 0$

then  $x = \mp y^{\frac{3}{2}} \Rightarrow \frac{dx}{dy} = \mp \frac{3}{2}y^{\frac{1}{2}}$



$$L = \int_0^1 \sqrt{1 + \frac{9}{4}y} dy + \int_0^4 \sqrt{1 + \frac{9}{4}y} dy = \frac{1}{18} \left[ \left. \frac{(4 + 9y)^{\frac{3}{2}}}{\frac{3}{2}} \right|_0^1 + \left. \frac{(4 + 9y)^{\frac{3}{2}}}{\frac{3}{2}} \right|_0^4 \right]$$

$$= \frac{1}{27} [(13\sqrt{13} - 4\sqrt{4}) + (40\sqrt{40} - 4\sqrt{4})] = 10.51$$

**EX-14** – Find the distance traveled between  $t=0$  and  $t=\frac{\pi}{2}$  a particle  $P(x,y)$  whose position at time  $t$  is given by:-  
 $x = a \cos t + a \cdot t \sin t$  and  $y = a \sin t - a \cdot t \cos t$  where  $a$  is a positive constant.

**Sol.**

$$x = a \cos t + a \cdot t \sin t \Rightarrow \frac{dx}{dt} = a \cdot t \cos t$$

$$y = a \sin t - a \cdot t \cos t \Rightarrow \frac{dy}{dt} = a \cdot t \sin t$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_0^{\frac{\pi}{2}} \sqrt{a^2 \cdot t^2 \cos^2 t + a^2 \cdot t^2 \sin^2 t} dt$$

$$= a \int_0^{\frac{\pi}{2}} t dt = \frac{a}{2} t^2 \Big|_0^{\frac{\pi}{2}} = \frac{a}{2} \left[ \frac{\pi^2}{4} - 0 \right] = \frac{a}{8} \pi^2$$

**EX-15** – Find the length of the curve:-

$$x = t - \sin t \text{ and } y = 1 - \cos t ; 0 \leq t \leq 2\pi$$

**Sol.**

$$x = t - \sin t \Rightarrow \frac{dx}{dt} = 1 - \cos t$$

$$y = 1 - \cos t \Rightarrow \frac{dy}{dt} = \sin t$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_0^{2\pi} \sqrt{(1 - \cos t)^2 + \sin^2 t} dt$$

$$= \int_0^{2\pi} \sqrt{1 - 2 \cos t + \cos^2 t + \sin^2 t} dt = \int_0^{2\pi} \sqrt{1 - 2 \cos t + 1} dt$$

$$= 2 \int_0^{2\pi} \sqrt{\frac{1 - \cos t}{2}} dt = 2 \int_0^{2\pi} \sin \frac{t}{2} dt = -4 \cos \frac{t}{2} \Big|_0^{2\pi}$$

$$= -4 [\cos \pi - \cos 0] = -4 [-1 - 1] = 8$$

### 7-5- The surface area:

Suppose that the curve  $y = f(x)$  is rotated about the  $x$ -axis. It will generate a surface in space. Then the surface area of the shape is:-

$$S = \int_a^b 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

If the curve rotated about the  $y$ -axis, then the surface area is:-

$$S = \int_c^d 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

If the curve sweeps out the surface is given in parametric form with  $x$  and  $y$  as functions of a third variable  $t$  that varies from  $t_a$  to  $t_b$  then we may compute the surface area from the formula:-

$$S = \int_{t_a}^{t_b} 2\pi \rho \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

where  $\rho$  is the distance from the axis of revolution to the element of arc length and is expressed as a function of  $t$ .

**EX-16** – The circle  $x^2 + y^2 = r^2$  is revolved about the  $x$ -axis. Find the area of the sphere generated.

**Sol.**

$$y = \sqrt{r^2 - x^2} \Rightarrow \frac{dy}{dx} = -\frac{x}{\sqrt{r^2 - x^2}}$$

$$\begin{aligned} S &= \int_a^b 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_{-r}^r 2\pi \sqrt{r^2 - x^2} \sqrt{1 + \frac{x^2}{r^2 - x^2}} dx = 2\pi r \int_{-r}^r dx \\ &= 2\pi r x \Big|_{-r}^r = 2\pi r(r - (-r)) = 4\pi r^2 \end{aligned}$$

**EX-17** – Find the area of the surface generated by rotating the portion of the curve  $y = \frac{1}{3}(x^2 + 2)^{\frac{3}{2}}$  between  $x=0$  and  $x=3$  about the  $y$ -axis.

**Sol.-**

$$y = \frac{1}{3}(x^2 + 2)^{\frac{3}{2}} \Rightarrow x = ((3y)^{\frac{2}{3}} - 2)^{\frac{1}{2}} \Rightarrow \frac{dx}{dy} = \frac{1}{(3y)^{\frac{1}{3}} \cdot ((3y)^{\frac{2}{3}} - 2)^{\frac{1}{2}}}$$

$$y = \frac{1}{3}(x^2 + 2)^{\frac{3}{2}} \quad \Rightarrow \Rightarrow \Rightarrow \quad \text{at } x=0 \quad y = \frac{2\sqrt{2}}{3} \quad \text{and} \quad \Rightarrow \Rightarrow \Rightarrow \quad \text{at } x=3 \quad y = \frac{11\sqrt{11}}{3}$$

$$S = \int_{\frac{2\sqrt{2}}{3}}^{\frac{11\sqrt{11}}{3}} 2\pi \sqrt{(3y)^{\frac{2}{3}} - 2} \cdot \sqrt{1 + \frac{1}{((3y)^{\frac{2}{3}} - 2)(3y)^{\frac{2}{3}}}} dy$$

$$= 2\pi \int_{\frac{2\sqrt{2}}{3}}^{\frac{11\sqrt{11}}{3}} \sqrt{\frac{(3y)^{\frac{4}{3}} - 2(3y)^{\frac{2}{3}} + 1}{(3y)^{\frac{2}{3}}}} dy = 2\pi \int_{\frac{2\sqrt{2}}{3}}^{\frac{11\sqrt{11}}{3}} \sqrt{\frac{((3y)^{\frac{2}{3}} - 1)^2}{(3y)^{\frac{2}{3}}}} dy$$

$$= 2\pi \int_{\frac{2\sqrt{2}}{3}}^{\frac{11\sqrt{11}}{3}} \left[ (3y)^{\frac{1}{3}} - (3y)^{-\frac{1}{3}} \right] dy = 2\pi \left[ \frac{1}{3} \frac{(3y)^{\frac{4}{3}}}{\frac{4}{3}} - \frac{1}{3} \frac{(3y)^{\frac{2}{3}}}{\frac{2}{3}} \right]_{\frac{2\sqrt{2}}{3}}^{\frac{11\sqrt{11}}{3}}$$

$$= \pi \left[ \frac{(3 \cdot \frac{11\sqrt{11}}{3})^{\frac{4}{3}}}{2} - (3 \cdot \frac{11\sqrt{11}}{3})^{\frac{2}{3}} - \frac{(3 \cdot \frac{2\sqrt{2}}{3})^{\frac{4}{3}}}{2} - (3 \cdot \frac{2\sqrt{2}}{3})^{\frac{2}{3}} \right] = \frac{99}{2} \pi$$

**EX-18** – The arc of the curve  $y = \frac{x^3}{3} + \frac{1}{4x}$  from  $x=1$  to  $x=3$  is rotated about the line  $y=-1$ . Find the surface area generated.

**Sol.-**



$$y = \frac{x^3}{3} + \frac{1}{4x} \Rightarrow \frac{dy}{dx} = x^2 - \frac{1}{4x^2} = \frac{4x^4 - 1}{4x^2}$$

$$\begin{aligned} S &= 2\pi \int_1^3 \left( \frac{x^3}{3} + \frac{1}{4x} + 1 \right) \sqrt{1 + \frac{(4x^4 - 1)^2}{16x^4}} dx \\ &= 2\pi \int_1^3 \frac{4x^4 + 12x + 3}{12x} \sqrt{\frac{(4x^4 + 1)^2}{16x^4}} dx \\ &= \frac{\pi}{24} \int_1^3 (16x^5 + 48x^2 + 16x + 12x^{-2} + 3x^{-3}) dx \\ &= \frac{\pi}{24} \left[ \frac{8}{3}x^6 + 16x^3 + 8x^2 - \frac{12}{x} - \frac{3}{2x^2} \right]_1^3 \\ &= \frac{\pi}{24} \left[ \frac{8}{3}(729 - 1) + 16(27 - 1) + 8(9 - 1) - 12\left(\frac{1}{3} - 1\right) - \frac{3}{2}\left(\frac{1}{9} - 1\right) \right] \\ &= \frac{1823}{18} \pi \end{aligned}$$

**EX-19** – Find the area of the surface generated by rotating the curve  $x = t^2$ ,  $y = t$ ,  $0 \leq t \leq 1$  about the x-axis.

**Sol.-**

$$x = t^2 \Rightarrow \frac{dx}{dt} = 2t \quad \text{and} \quad y = t \Rightarrow \frac{dy}{dt} = 1$$

$$\begin{aligned} S &= \int_{t_a}^{t_b} 2\pi \rho \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = 2\pi \int_0^1 t \sqrt{4t^2 + 1} dt \\ &= \frac{\pi}{4} \left[ \frac{(4t^2 + 1)^{\frac{3}{2}}}{\frac{3}{2}} \right]_0^1 = \frac{\pi}{6} [5\sqrt{5} - 1] \end{aligned}$$

## Problems – 7

1) Find the area of the region bounded by the given curves and lines for the following problems:-

1. The coordinate axes and the line  $x + y = a$
2. The  $x$ -axis and the curve  $y = e^x$  and the lines  $x = 0$  ,  $x = 1$
3. The curve  $y^2 + x = 0$  and the line  $y = x + 2$
4. The curves  $x = y^2$  and  $x = 2y - y^2$
5. The parabola  $x = y - y^2$  and the line  $x + y = 0$

$$(ans. : 1. \frac{a^2}{2} ; 2. e - 1 ; 3. \frac{9}{2} ; 4. \frac{1}{3} ; 5. \frac{4}{3})$$

2) Write an equivalent double integral with order of integration reversed for each integrals check your answer by evaluation both double integrals, and sketch the region.

$$1. \int_0^2 \int_1^{e^x} dy dx \qquad (ans. : \int_1^2 \int_{\ln y}^2 dx dy ; e^2 - 3)$$

$$2. \int_0^1 \int_{\sqrt{y}}^1 dx dy \qquad (ans. : \int_0^1 \int_0^{x^2} dy dx ; \frac{1}{3})$$

$$3. \int_0^{\sqrt{2}} \int_{-\sqrt{4-2y^2}}^{\sqrt{4-2y^2}} y dx dy \qquad (ans. : \int_{-2}^2 \int_0^{\sqrt{4-x^2}} y dy dx ; \frac{8}{3})$$

3) Find the volume of the tetrahedron bounded by the plane  $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$  and the coordinate planes.

$$(ans. : \frac{1}{6} |abc|)$$

4) Find the volume bounded by the plane  $z = 0$  laterally by the elliptic cylinder  $x^2 + 4y^2 = 4$  and above by the plane  $z = x + 2$ .

$$(ans. : 4\pi)$$

5) Find the lengths of the following curves:-

1.  $y = x^{\frac{3}{2}}$  from  $(0,0)$  to  $(4,8)$  (ans. :  $\frac{8}{27}(10\sqrt{10} - 1)$ )

2.  $y = \frac{x^3}{3} + \frac{1}{4x}$  from  $x = 1$  to  $x = 3$  (ans. :  $\frac{53}{6}$ )

3.  $x = \frac{y^4}{4} + \frac{1}{8y^2}$  from  $y = 1$  to  $y = 2$  (ans. :  $\frac{123}{32}$ )

4.  $(y + 1)^2 = 4x^3$  from  $x = 0$  to  $x = 1$  (ans. :  $\frac{4}{27}(10\sqrt{10} - 1)$ )

6) Find the distance traveled by the particle  $P(x,y)$  between  $t=0$  and  $t=4$  if the position at time  $t$  is given by:  $x = \frac{t^2}{2}$  ;  $y = \frac{1}{3}(2t + 1)^{\frac{3}{2}}$   
(ans. : 12)

7) The position of a particle  $P(x,y)$  at time  $t$  is given by:  
 $x = \frac{1}{3}(2t + 3)^{\frac{3}{2}}$  ;  $y = \frac{t^2}{2} + t$  . Find the distance it travel between  $t=0$   
and  $t=3$ . (ans. :  $\frac{21}{2}$ )

8) Find the area of the surface generated by rotating about the x-axis  
the arc of the curve  $y = x^3$  between  $x = 0$  and  $x = 1$ .  
(ans. :  $\frac{\pi}{27}(10\sqrt{10} - 1)$ )

9) Find the area of the surface generated by rotating about the y-axis  
the arc of the curve  $y = x^2$  between  $(0,0)$  and  $(2,4)$  .  
(ans. :  $\frac{\pi}{6}(17\sqrt{17} - 1)$ )

10) Find the area of the surface generated by rotating about the y-  
axis the curve  $y = \frac{x^2}{2} + \frac{1}{2}$  ;  $0 \leq x \leq 1$  . (ans. :  $\frac{2}{3}\pi(2\sqrt{2} - 1)$ )

11) The curve described by the particle  $P(x,y)$   $x = t + 1$  ,  $y = \frac{t^2}{2} + t$   
from  $t = 0$  to  $t = 4$  is rotated about the y-axis. Find the surface area  
that is generated.

(ans. :  $\frac{2\sqrt{2}}{3}\pi(13\sqrt{13} - 1)$ )