


# Blood Clotting



**Prepared and Presented by:**

**Lecturer Dr/ Ayad AbdElSalam  
Assist. Lecturer Dr/ Ghadeer Talib**



**Teaching of Physiology  
College of Technology & Health Sciences  
Radiological Techniques Department**

# **THROMBOCYTES (Platelets)**

- Small, irregularly-shaped, oval non-nucleated cells, 2-4  $\mu\text{m}$  in diameter.
- Derived from fragmentation of precursor megakaryocytes (bone marrow).
- Reference range (SI units) is 150,000 - 450,000  $\mu\text{L}$ .
- The average lifespan of a platelet is between 8 and 12 days.
- Old platelets are destroyed by phagocytosis in the spleen and by Kupffer cells in the liver.

## **Platelet functions:**

- Maintenance of Vascular Integrity.
- Formation of the Primary Hemostatic Plug.
- Accelerate thrombin formation.




# HEMOSTASIS (blood clots)

Blood coagulation or clotting is an important phenomenon to prevent excess loss of blood in case of injury or trauma. The blood stops flowing from a wound in case of injury. The blood clot or 'coagulum' is formed by a network of fibrin threads.

## HEMOSTASIS


- 1) Constriction of the blood vessel.
- 2) Formation of a temporary "platelet plug."
- 3) Activation of the coagulation cascade.
- 4) Formation of "fibrin plug" or the final clot.



**Prothrombin** is the inactive form of thrombin that is present in the plasma. Thrombokinase converts prothrombin to active thrombin which in turn activates fibrinogen to fibrin. All these clotting factors help in blood coagulation.

An injury stimulates platelets or thrombocytes to release various factors that initiate the blood clotting cascade. Calcium ions play an important role in blood coagulation.

## Human Blood Clotting



## **Primary hemostasis (platelet clotting)**

**Primary hemostasis is when your body forms a temporary plug to seal an injury. To accomplish that, platelets that circulate in your blood stick to the damaged tissue and activate. That activation means they can “recruit” more platelets to form a platelet “plug” to stop blood loss from the damaged area. That clot works much like a cork or bottle stopper, keeping blood in and debris or germs out.**

**Primary hemostasis may also involve constriction (narrowing) of the damaged blood vessel, which can happen because of substances that activated platelets release.**



## **Secondary hemostasis (coagulation cascade)**

**The platelet plug is the first step to stop bleeding, but it isn't stable enough to stay in place without help. The next step, which stabilizes the plug, is secondary hemostasis. This step, sometimes called coagulation, involves molecules in your blood called "coagulation factors." Those factors activate in sequence, the "coagulation cascade," which amplifies clotting effects as the sequence continues. Ultimately, the coagulation cascade forms a substance called fibrin. During this step, the platelet plug acts like bricks and the fibrin acts like mortar. Together, they form a solid, stable clot.**

**Coagulation of blood occurs through a series of reactions due to the activation of a group of substances called clotting factors.**

**There are 13 clotting factors identified and named after the scientists who discovered them or as per the activity. Only factor IX or Christmas factor is named after the patient in whom it was discovered.**

<b>Factor I Fibrinogen</b>	<b>Factor VI Unassigned</b>	<b>Factor XI Plasma thromboplastin antecedent</b>
<b>Factor II Prothrombin</b>	<b>Factor VII Stable factor</b>	<b>Factor XII Hageman factor</b>
<b>Factor III Thromboplastin</b>	<b>Factor VIII Antihemophilic factor</b>	<b>Factor XIII Fibrin-stabilising factor</b>
<b>Factor IV Calcium Factor</b>	<b>Factor IX Christmas factor</b>	
<b>Factor V Labile factor (Proaccelerin)</b>	<b>Factor X Stuart-Prower factor</b>	

**Prothrombin activators are a group of substances which convert prothrombin to thrombin in two ways:**


**1. Extrinsic pathway: (the main pathway to initiate coagulation) The process is started when injured endothelial cells produce tissue factor (factor III), which activates factor VII.**

**2. Intrinsic pathway: (which amplifies coagulation) involves the activation of factors XII, XI, IX, and factor VIII.**

**The prothrombin activator converts prothrombin to thrombin in the presence of enough ionic  $\text{Ca}^{++}$  from platelets.**

**The most significant ions in the process of coagulation of blood are calcium ions. Because calcium ions are required for the conversion of thromboplastin into prothrombinase and prothrombin into thrombin (active).**

# Blood coagulation *in vivo*



## **Fibrin clot remodeling**

**The last stage of hemostasis is when your body remodels the existing clot into a fibrin clot. Your body does that because blood clots are a temporary patch, not a permanent solution. That removal involves a process called fibrinolysis. During fibrinolysis, your body remodels the clot into the same kind of tissue that was there before the injury.**

**The two final phases in the hemostatic process, plasma coagulation with the formation of a fibrin clot, and fibrinolysis leading to the dissolution of fibrin clots**