

1st Semester (2023-2024)

Lecture No. 3

Derivatives

Let $y = f(x)$ be a function of x . If the limit :

$$\frac{dy}{dx} = f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$$

exists and is finite , we call this limit the derivative of f at x and say that f is differentiable at x .

EX-1 – Find the derivative of the function : $f(x) = \frac{1}{\sqrt{2x+3}}$

Sol.:

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{\sqrt{2(x + \Delta x) + 3}} - \frac{1}{\sqrt{2x + 3}}}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{2x + 3} - \sqrt{2(x + \Delta x) + 3}}{\Delta x \cdot \sqrt{2(x + \Delta x) + 3} \sqrt{2x + 3}} \cdot \frac{\sqrt{2x + 3} + \sqrt{2(x + \Delta x) + 3}}{\sqrt{2x + 3} + \sqrt{2(x + \Delta x) + 3}} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(2x + 3) - (2(x + \Delta x) + 3)}{\Delta x \cdot \sqrt{2(x + \Delta x) + 3} \sqrt{2x + 3} (\sqrt{2x + 3} + \sqrt{2(x + \Delta x) + 3})} \\ &= \frac{-2}{(2x + 3)(\sqrt{2x + 3} + \sqrt{2x + 3})} = -\frac{1}{\sqrt{(2x + 3)^3}} \end{aligned}$$

Rules of derivatives : Let c and n are constants, u , v and w are differentiable functions of x :

1. $\frac{d}{dx} c = 0$
2. $\frac{d}{dx} u^n = n u^{n-1} \frac{du}{dx} \Rightarrow \frac{d}{dx} \left(\frac{1}{u} \right) = -\frac{1}{u^2} \frac{du}{dx}$
3. $\frac{d}{dx} cu = c \frac{du}{dx}$
4. $\frac{d}{dx} (u \mp v) = \frac{du}{dx} \mp \frac{dv}{dx} ; \frac{d}{dx} (u \mp v \mp w) = \frac{du}{dx} \mp \frac{dv}{dx} \mp \frac{dw}{dx}$
5. $\frac{d}{dx} (u \cdot v) = u \cdot \frac{dv}{dx} + v \frac{du}{dx}$

1st Semester (2023-2024)

and $\frac{d}{dx}(u.v.w) = u.v \frac{dw}{dx} + u.w \frac{dv}{dx} + v.w \frac{du}{dx}$

6. $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$ where $v \neq 0$

EX-2- Find $\frac{dy}{dx}$ for the following functions :

a) $y = (x^2 + 1)^5$

b) $y = [(5-x)(4-2x)]^2$

c) $y = (2x^3 - 3x^2 + 6x)^{-5}$

d) $y = \frac{12}{x} - \frac{4}{x^3} + \frac{3}{x^4}$

e) $y = \frac{(x^2 + x)(x^2 - x + 1)}{x^3}$

f) $y = \frac{x^2 - 1}{x^2 + x - 2}$

Sol.-

a) $\frac{dy}{dx} = 5(x^2 + 1)^4 \cdot 2x = 10x(x^2 + 1)^4$

b) $\frac{dy}{dx} = 2[(5-x)(4-2x)][-2(5-x) - (4-2x)]$
 $= 8(5-x)(2-x)(2x-7)$

c) $\frac{dy}{dx} = -5(2x^3 - 3x^2 + 6x)^{-6}(6x^2 - 6x + 6)$
 $= -30(2x^3 - 3x^2 + 6x)^{-6}(x^2 - x + 1)$

d) $y = 12x^{-1} - 4x^{-3} + 3x^{-4} \Rightarrow \frac{dy}{dx} = -12x^{-2} + 12x^{-4} - 12x^{-5}$
 $\Rightarrow \frac{dy}{dx} = -\frac{12}{x^2} + \frac{12}{x^4} - \frac{12}{x^5}$

e) $y = \frac{(x+1)(x^2 - x + 1)}{x^3} \Rightarrow$
 $\frac{dy}{dx} = \frac{x^3[(x^2 - x + 1) + (x+1)(2x-1)] - 3x^2(x+1)(x^2 - x + 1)}{x^6} = -\frac{3}{x^4}$

f) $\frac{dy}{dx} = \frac{2x(x^2 + x - 2) - (x^2 - 1)(2x+1)}{(x^2 + x - 2)^2} = \frac{x^2 - 2x + 1}{(x^2 + x - 2)^2}$

1st Semester (2023-2024)

The Chain Rule:

1. Suppose that $h = g \circ f$ is the composite of the differentiable functions $y = g(t)$ and $x = f(t)$, then h is a differentiable function of x whose derivative at each value of x is :

$$\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$$

2. If y is a differentiable function of t and t is differentiable function of x , then y is a differentiable function of x :

$$y = g(t) \text{ and } t = f(x) \Rightarrow \frac{dy}{dx} = \frac{dy}{dt} * \frac{dt}{dx}$$

EX-3 – Use the chain rule to express dy/dx in terms of x and y :

- a) $y = \frac{t^2}{t^2 + 1}$ and $t = \sqrt{2x + 1}$
 b) $y = \frac{1}{t^2 + 1}$ and $x = \sqrt{4t + 1}$
 c) $y = \left(\frac{t-1}{t+1}\right)^2$ and $x = \frac{1}{t^2} - 1$ at $t = 2$
 d) $y = 1 - \frac{1}{t}$ and $t = \frac{1}{1-x}$ at $x = 2$

Sol.-

$$\begin{aligned}
 a) \quad & y = \frac{t^2}{t^2 + 1} \Rightarrow \frac{dy}{dt} = \frac{2t(t^2 + 1) - 2t \cdot t^2}{(t^2 + 1)^2} = \frac{2t}{(t^2 + 1)^2} \\
 & t = (2x + 1)^{\frac{1}{2}} \Rightarrow \frac{dt}{dx} = \frac{1}{2} \cdot (2x + 1)^{-\frac{1}{2}} \cdot 2 = \frac{1}{\sqrt{2x + 1}} \\
 & \frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = \frac{2t}{(t^2 + 1)^2} \cdot \frac{1}{\sqrt{2x + 1}} = \frac{2\sqrt{2x + 1}}{((2x + 1) + 1)^2} \cdot \frac{1}{\sqrt{2x + 1}} = \frac{1}{2(x + 1)^2}
 \end{aligned}$$

1st Semester (2023-2024)

$$b) \quad y = (t^2 + 1)^{-1} \Rightarrow \frac{dy}{dx} = -2t(t^2 + 1)^{-2} = -\frac{2t}{(t^2 + 1)^2}$$

$$x = (4t + 1)^{\frac{1}{2}} \Rightarrow \frac{dx}{dt} = \frac{1}{2}(4t + 1)^{-\frac{1}{2}} \cdot 4 = \frac{2}{\sqrt{4t + 1}}$$

$$\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt} = -\frac{2t}{(t^2 + 1)^2} \div \frac{2}{\sqrt{4t + 1}} = -\frac{t\sqrt{4t + 1}}{(t^2 + 1)^2}$$

$$= -\frac{x^2 - 1}{4} \cdot x \div \frac{1}{y^2} = -\frac{xy^2(x^2 - 1)}{4}$$

$$\text{where } x = \sqrt{4t + 1} \Rightarrow t = \frac{x^2 - 1}{4}$$

$$\text{where } y = \frac{1}{t^2 + 1} \Rightarrow t^2 + 1 = \frac{1}{y}$$

$$c) \quad y = \left(\frac{t-1}{t+1}\right)^2 \Rightarrow \frac{dy}{dt} = 2\left(\frac{t-1}{t+1}\right) \frac{t+1-(t-1)}{(t+1)^2} = \frac{4(t-1)}{(t+1)^3}$$

$$\Rightarrow \left[\frac{dy}{dt} \right]_{t=2} = \frac{4(2-1)}{(2+1)^3} = \frac{4}{27}$$

$$x = \frac{1}{t^2} - 1 \Rightarrow \frac{dx}{dt} = -\frac{2}{t^3} \Rightarrow \left[\frac{dx}{dt} \right]_{t=2} = -\frac{2}{2^3} = -\frac{1}{4}$$

$$\left[\frac{dy}{dx} \right]_{t=2} = \left[\frac{dy}{dt} \div \frac{dx}{dt} \right]_{t=2} = \frac{4}{27} \div \left(-\frac{1}{4} \right) = -\frac{16}{27}$$

$$d) \quad t = \frac{1}{1-x} = \frac{1}{1-2} = -1 \quad \text{at } x = 2$$

$$y = 1 - \frac{1}{t} \Rightarrow \frac{dy}{dt} = \frac{1}{t^2} \Rightarrow \left[\frac{dy}{dt} \right]_{t=-1} = \frac{1}{(-1)^2} = 1$$

$$t = (1-x)^{-1} \Rightarrow \frac{dt}{dx} = -(1-x)^{-2}(-1) = \frac{1}{(1-x)^2}$$

$$\Rightarrow \left[\frac{dt}{dx} \right]_{x=2} = \frac{1}{(1-2)^2} = 1$$

$$\left[\frac{dy}{dx} \right]_{x=2} = \left[\frac{dy}{dt} \right]_{x=2} \cdot \left[\frac{dt}{dx} \right]_{x=2} = 1 * 1 = 1$$

Higher derivatives : If a function $y = f(x)$ possesses a derivative at every point of some interval , we may form the function $f'(x)$ and talk

1st Semester (2023-2024)

about its derivate , if it has one . The procedure is formally identical with that used before , that is :

$$\frac{d^2 y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f'(x + \Delta x) - f'(x)}{\Delta x}$$

if the limit exists .

This derivative is called the second derivative of y with respect to x . It is written in a number of ways , for example,

$$y'', f''(x), \text{ or } \frac{d^2 f(x)}{dx^2}.$$

In the same manner we may define third and higher derivatives , using similar notations . The n th derivative may be written :

$$y^{(n)}, f^{(n)}(x), \frac{d^n y}{dx^n} .$$

EX-4- Find all derivatives of the following function :

$$y = 3x^3 - 4x^2 + 7x + 10$$

Sol.-

$$\begin{aligned} \frac{dy}{dx} &= 9x^2 - 8x + 7 & , \quad \frac{d^2 y}{dx^2} &= 18x - 8 \\ \frac{d^3 y}{dx^3} &= 18 & , \quad \frac{d^4 y}{dx^4} &= 0 = \frac{d^5 y}{dx^5} = \dots \end{aligned}$$

Ex-5 – Find the third derivative of the following function :

$$y = \frac{1}{x} + \sqrt{x^3}$$

Sol.-

$$\begin{aligned} \frac{dy}{dx} &= -\frac{1}{x^2} + \frac{3}{2} x^{\frac{1}{2}} \\ \frac{d^2 y}{dx^2} &= \frac{2}{x^3} + \frac{3}{4} x^{-\frac{1}{2}} \\ \frac{d^3 y}{dx^3} &= -\frac{6}{x^4} - \frac{3}{8} x^{-\frac{3}{2}} \Rightarrow \frac{d^3 y}{dx^3} = -\frac{6}{x^4} - \frac{3}{8\sqrt{x^3}} \end{aligned}$$

1st Semester (2023-2024)

Implicit Differentiation: If the formula for f is an algebraic combination of powers of x and y . To calculate the derivatives of these implicitly defined functions , we simply differentiate both sides of the defining equation with respect to x .

EX-6- Find $\frac{dy}{dx}$ for the following functions:

$$a) x^2 \cdot y^2 = x^2 + y^2 \quad b) (x+y)^3 + (x-y)^3 = x^4 + y^4$$

$$c) \frac{x-y}{x-2y} = 2 \text{ at } P(3,1) \quad d) xy + 2x - 5y = 2 \text{ at } P(3,2)$$

Sol.

$$a) x^2(2y\frac{dy}{dx}) + y^2(2x) = 2x + 2y\frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{x - xy^2}{x^2y - y}$$

$$b) 3(x+y)^2(1+\frac{dy}{dx}) + 3(x-y)^2(1-\frac{dy}{dx}) = 4x^3 + 4y^3 \frac{dy}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{4x^3 - 3(x+y)^2 - 3(x-y)^2}{3(x+y)^2 - 3(x-y)^2 - 4y^3} \Rightarrow \frac{dy}{dx} = \frac{2x^3 - 3x^2 - 3y^2}{6xy - 2y^3}$$

$$c) \frac{(x-2y)(1-\frac{dy}{dx}) - (x-y)(1-2\frac{dy}{dx})}{(x-2y)^2} = 0 \Rightarrow \frac{dy}{dx} = \frac{y}{x} \Rightarrow \left[\frac{dy}{dx} \right]_{(3,1)} = \frac{1}{3}$$

$$d) x\frac{dy}{dx} + y + 2 - 5\frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = \frac{y+2}{5-x} \Rightarrow \left[\frac{dy}{dx} \right]_{(3,2)} = \frac{2+2}{5-3} = 2$$

Exponential functions: If u is any differentiable function of x , then :

$$7) \frac{d}{dx} a^u = a^u \cdot \ln a \cdot \frac{du}{dx} \quad \text{and} \quad \frac{d}{dx} e^u = e^u \cdot \frac{du}{dx}$$

1st Semester (2023-2024)

EX-7 –Find $\frac{dy}{dx}$ for the following functions :

a) $y = 2^{3x}$

b) $y = 2^x \cdot 3^x$

c) $y = (2^x)^2$

d) $y = x \cdot 2^{x^2}$

e) $y = e^{(x+e^{5x})}$

f) $y = e^{\sqrt{1+5x^2}}$

Sol.-

a) $y = 2^{3x} \Rightarrow \frac{dy}{dx} = 2^{3x} \cdot 3 \ln 2$

b) $y = 2^x \cdot 3^x \Rightarrow y = 6^x \Rightarrow \frac{dy}{dx} = 6^x \cdot \ln 6$

c) $y = (2^x)^2 \Rightarrow y = 2^{2x} \Rightarrow \frac{dy}{dx} = 2^{2x} \ln 2 \cdot 2 = 2^{2x+1} \ln 2$

d) $y = x \cdot 2^{x^2} \Rightarrow \frac{dy}{dx} = x \cdot 2^{x^2} \ln 2 \cdot 2x + 2^{x^2} = 2^{x^2} (2x^2 \ln 2 + 1)$

e) $y = e^{(x+e^{5x})} \Rightarrow \frac{dy}{dx} = e^{(x+e^{5x})} (1 + 5e^{5x})$

f) $y = e^{(1+5x^2)^{\frac{1}{2}}} \Rightarrow \frac{dy}{dx} = e^{(1+5x^2)^{\frac{1}{2}}} \frac{1}{2} (1+5x^2)^{-\frac{1}{2}} \cdot 10x = e^{\sqrt{1+5x^2}} \frac{5x}{\sqrt{1+5x^2}}$

Logarithm functions : If u is any differentiable function of x , then :

8) $\frac{d}{dx} \log_a u = \frac{1}{u \cdot \ln a} \cdot \frac{du}{dx}$ and $\frac{d}{dx} \ln u = \frac{1}{u} \cdot \frac{du}{dx}$

EX-8 –Find $\frac{dy}{dx}$ for the following functions :

a) $y = \log_{10} e^x$

b) $y = \log_5 (x+1)^2$

c) $y = \log_2 (3x^2 + 1)^3$

d) $y = [\ln(x^2 + 2)^2]^3$

e) $y + \ln(xy) = 1$

f) $y = \frac{(2x^3 - 4)^{\frac{2}{3}} \cdot (2x^2 + 3)^{\frac{5}{2}}}{(7x^3 + 4x - 3)^2}$

Sol. –

1st Semester (2023-2024)

$$a) \ y = \log_{10} e^x \Rightarrow y = x \log_{10} e \Rightarrow \frac{dy}{dx} = \log_{10} e = \frac{\ln e}{\ln 10} = \frac{1}{\ln 10}$$

$$b) \ y = \log_5 (x+1)^2 = 2 \log_5 (x+1) \Rightarrow \frac{dy}{dx} = \frac{2}{(x+1)\ln 5}$$

$$c) \ y = 3 \log_2 (3x^2 + 1) \Rightarrow \frac{dy}{dx} = \frac{3}{3x^2 + 1} \cdot \frac{6x}{\ln 2} = \frac{18x}{(3x^2 + 1)\ln 2}$$

$$d) \ \frac{dy}{dx} = 3[2 \ln(x^2 + 2)]^2 \frac{2}{x^2 + 2} \cdot 2x = \frac{48x[\ln(x^2 + 2)]^2}{x^2 + 2}$$

$$e) \ y + \ln x + \ln y = 1 \Rightarrow \frac{dy}{dx} + \frac{1}{x} + \frac{1}{y} \cdot \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{y}{x(y+1)}$$

$$f) \ \ln y = \frac{2}{3} \ln(2x^3 - 4) + \frac{5}{2} \ln(2x^2 + 3) - 2 \ln(7x^3 + 4x - 3) \\ \Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \frac{2}{3} \cdot \frac{6x^2}{2x^3 - 4} + \frac{5}{2} \cdot \frac{4x}{2x^2 + 3} - 2 \cdot \frac{21x^2 + 4}{7x^3 + 4x - 3} \\ \Rightarrow \frac{dy}{dx} = 2y \left[\frac{2x^2}{2x^3 - 4} + \frac{5x}{2x^2 + 3} - \frac{21x^2 + 4}{7x^3 + 4x - 3} \right]$$

Trigonometric functions : If u is any differentiable function of x , then :

$$9) \ \frac{d}{dx} \sin u = \cos u \cdot \frac{du}{dx}$$

$$10) \ \frac{d}{dx} \cos u = -\sin u \cdot \frac{du}{dx}$$

$$11) \ \frac{d}{dx} \tan u = \sec^2 u \cdot \frac{du}{dx}$$

$$12) \ \frac{d}{dx} \cot u = -\csc^2 u \cdot \frac{du}{dx}$$

$$13) \ \frac{d}{dx} \sec u = \sec u \cdot \tan u \cdot \frac{du}{dx}$$

$$14) \ \frac{d}{dx} \csc u = -\csc u \cdot \cot u \cdot \frac{du}{dx}$$

EX-9- Find $\frac{dy}{dx}$ for the following functions :

1st Semester (2023-2024)

- | | |
|--|--|
| <i>a) $y = \tan(3x^2)$</i> | <i>b) $y = (\csc x + \cot x)^2$</i> |
| <i>c) $y = 2\sin\frac{x}{2} - x\cos\frac{x}{2}$</i> | <i>d) $y = \tan^2(\cos x)$</i> |
| <i>e) $x + \tan(xy) = 0$</i> | <i>f) $y = \sec^4 x - \tan^4 x$</i> |

Sol.-

$$\begin{aligned}
 a) \frac{dy}{dx} &= \sec^2(3x^2) \cdot 6x = 6x \cdot \sec^2(3x^2) \\
 b) \frac{dy}{dx} &= 2(\csc x + \cot x)(-\csc x \cdot \cot x - \csc^2 x) = -2 \csc x \cdot (\csc x + \cot x)^2 \\
 c) \frac{dy}{dx} &= 2 \cos \frac{x}{2} \cdot \frac{1}{2} - \left[x(-\sin \frac{x}{2}) \cdot \frac{1}{2} + \cos \frac{x}{2} \right] = \frac{x}{2} \cdot \sin \frac{x}{2} \\
 d) \frac{dy}{dx} &= 2 \cdot \tan(\cos x) \cdot \sec^2(\cos x) \cdot (-\sin x) = -2 \cdot \sin x \cdot \tan(\cos x) \cdot \sec^2(\cos x) \\
 e) 1 + \sec^2(xy) \cdot (x \frac{dy}{dx} + y) &= 0 \Rightarrow \frac{dy}{dx} = -\frac{1 + y \cdot \sec^2(xy)}{x \cdot \sec^2(xy)} = -\frac{\cos^2(xy) + y}{x} \\
 f) \frac{dy}{dx} &= 4 \sec^3 x \cdot \sec x \cdot \tan x - 4 \cdot \tan^3 x \cdot \sec^2 x = 4 \tan x \cdot \sec^2 x
 \end{aligned}$$

EX-10- Prove that :

$$a) \frac{d}{dx} \tan u = \sec^2 u \cdot \frac{du}{dx} \quad b) \frac{d}{dx} \sec u = \sec u \cdot \tan u \cdot \frac{du}{dx}$$

Proof:

$$\begin{aligned}
 a) L.H.S. &= \frac{d}{dx} \tan u = \frac{d}{dx} \frac{\sin u}{\cos u} = \frac{\cos u \cdot \cos u \cdot \frac{du}{dx} - \sin u \cdot (-\sin u) \frac{du}{dx}}{\cos^2 u} \\
 &= \frac{\cos^2 u + \sin^2 u}{\cos^2 u} \cdot \frac{du}{dx} = \frac{1}{\cos^2 u} \cdot \frac{du}{dx} = \sec^2 u \cdot \frac{du}{dx} = R.H.S.
 \end{aligned}$$

$$\begin{aligned}
 b) L.H.S. &= \frac{d}{dx} \sec u = \frac{d}{dx} \frac{1}{\cos u} = -\frac{1}{\cos^2 u} (-\sin u) \frac{du}{dx} \\
 &= \frac{1}{\cos u} \cdot \frac{\sin u}{\cos u} \cdot \frac{du}{dx} = \sec u \cdot \tan u \cdot \frac{du}{dx} = R.H.S.
 \end{aligned}$$

The inverse trigonometric functions : If u is any differentiable function

1st Semester (2023-2024)

of x , then :

$$15) \frac{d}{dx} \sin^{-1} u = \frac{1}{\sqrt{1-u^2}} \frac{du}{dx} \quad -1 < u < 1$$

$$16) \frac{d}{dx} \cos^{-1} u = -\frac{1}{\sqrt{1-u^2}} \frac{du}{dx} \quad -1 < u < 1$$

$$17) \frac{d}{dx} \tan^{-1} u = \frac{1}{1+u^2} \frac{du}{dx}$$

$$18) \frac{d}{dx} \cot^{-1} u = -\frac{1}{1+u^2} \frac{du}{dx}$$

$$19) \frac{d}{dx} \sec^{-1} u = \frac{1}{|u|\sqrt{u^2-1}} \frac{du}{dx} \quad |u| > 1$$

$$20) \frac{d}{dx} \csc^{-1} u = -\frac{1}{|u|\sqrt{u^2-1}} \frac{du}{dx} \quad |u| > 1$$

EX-11- Find $\frac{dy}{dx}$ in each of the following functions :

$$a) y = \cot^{-1} \frac{2}{x} + \tan^{-1} \frac{x}{2}$$

$$b) y = \sin^{-1} \frac{x-1}{x+1}$$

$$c) y = x \cdot \cos^{-1} 2x - \frac{1}{2} \sqrt{1-4x^2}$$

$$d) y = \sec^{-1} 5x$$

$$e) y = x \cdot \ln(\sec^{-1} x)$$

$$f) y = 3^{\sin^{-1} 2x}$$

Sol. –

$$a) \frac{dy}{dx} = -\frac{1}{1+\left(\frac{2}{x}\right)^2} \cdot 2 \left(-\frac{1}{x^2}\right) + \frac{1}{1+\left(\frac{x}{2}\right)^2} \cdot \frac{1}{2} = \frac{4}{4+x^2}$$

$$b) \frac{dy}{dx} = \frac{1}{\sqrt{1-\left(\frac{x-1}{x+1}\right)^2}} \cdot \frac{(x+1).1-(x-1).1}{(x+1)^2} = \frac{1}{(x+1)\sqrt{x}}$$

$$c) \frac{dy}{dx} = x \frac{-2}{\sqrt{1-4x^2}} + \cos^{-1} 2x - \frac{1}{4} \cdot \frac{-8x}{\sqrt{1-4x^2}} = \cos^{-1} 2x$$

$$d) \frac{dy}{dx} = \frac{5}{|5x|\sqrt{25x^2-1}} = \frac{1}{|x|\sqrt{25x^2-1}}$$