

وزارة التعليم و البحث العلمي
جامعة المستقبل
قسم الفيزياء الطبية
المرحلة الاولى/ مختبر الميكانيك II

تجربة قانون بويل
Boyle's Law experiment

اعداد : م . زهراء عباس علي

The aim of Experiment: Boyle's Law Investigation and Measure the pressure of the atmosphere.

Apparatus: 1-A ruler of metric scale . (100cm)

2-Glass tube connected with a closed plastic tube at the end .

3-Liquid mercury (Hg).

□Theory:

*Boyle's law states that, for constant temperature, the product of the volume and the pressure of an ideal gas is a constant.

$$*PV=C\text{..... (1)}$$

$$*The\ ideal\ gas\ law\ PV=nRT\ \text{..... (2)}$$

*states that this constant (nRT) is proportional to the amount of ideal gas in the sample (the number of moles, n).

*The absolute temperature, T.

*The constant R in this equation is the universal gas constant which has a value of $R=8.31\ J/(mole.K)$ in SI unit.

*Note that if T is held constant throughout the experiment, then the ideal gas law reduces to Boyle's law.

*An experiment to investigate Boyle's law is carried out with the apparatus shown in the diagram.

*The pressure and volume of the gas (air) trapped in the closed end can be varied by raising or lowering the other end.

*By measuring the difference in levels of mercury in the two tubes the pressure of the gas in the closed end can be calculated.

*The volume of gas in this end can be calculated by assuming that the glass tube is a cylinder.

Diagram of Experiment:




Table of reading :

h(cm)	L cm	1/L (cm ⁻¹)	P=ρgh	V(cm ³)
0	12			
5	11.5			
10	11			
15	10.5			
20	10			

Results and your calculation :

$$P = \rho gh \dots (1)$$

ρ = density of Mercury (Hg) = 13600 Kg/m³

1mm.Hg = 132.3 pa .

$g = 9.8 \text{ m/sec}^2$ or $= 10 \text{ m/sec}^2$

$$v = 4\pi^2 L$$