



**Ministry of Higher Education and Scientific Research**  
**Al-Mustaqbal University College**  
**Department of Chemical Engineering and petroleum**  
**Industrials**

**Week: 4,5**

**Mathematics II**

**2<sup>nd</sup> Stage**

**Lecturer: Sara I. Mohammed**

2019-2020

**Triple integral**


If  $f(x, y, z)$  is a function defined on a closed bounded region  $D$  in space, such as the region occupied by a solid ball or a lump of clay, then the integral of  $f$  over  $D$  may be defined in the following way.

$$V = \iiint_D dV = \int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} \int_{z=f_1(x,y)}^{z=f_2(x,y)} F(x, y, z) dz dy dx$$

**a- Surface area**

Let  $f(x, y)$  be a differentiable function. As we have seen,  $z=f(x, y)$  defines a surface in  $x, y, z$ -space. In some applications, it is necessary to know the surface area of the surface above some region  $R$  in the  $xy$ -plane. See the figure.

$$S = \iint_R \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dy dx$$





**Examples**


**1. Double integral**

**a- Cartesian form**


1- Find the limits of the following integral




(a)



(b)



(c)



(d)

(e)

$$(a) \int_0^9 \int_0^{\sqrt{x}} dy dx$$

$$\int_0^3 \int_{y^2}^9 dx dy$$

$$\int_0^{\pi/4} \int_{\tan x}^1 dy dx$$

$$\int_0^1 \int_0^{\tan^{-1} y} dx dy$$

$$(e) \int_0^1 \int_x^{3-2x} dy dx$$

$$\int_0^1 \int_0^y dx dy + \int_1^3 \int_0^{(3-y)/2} dx dy$$

$$\int_0^1 \int_0^1 dy dx + \int_1^e \int_{\ln x}^1 dy dx = \int_{-1}^2 \int_{x^2}^{x+2} dy dx$$

$$\int_0^1 \int_0^{e^y} dx dy = \int_0^1 \int_{-\sqrt{y}}^{\sqrt{y}} dx dy + \int_1^3 \int_{y-2}^{\sqrt{y}} dx dy$$

(f)

(g)

4

2- Evaluate the following

a-  $\int_0^3 \int_1^2 (1 + 8xy) \, dy \, dx$        $\int_0^1 \int_0^1 \left(1 - \frac{x^2 + y^2}{2}\right) \, dx \, dy$

c-  $\int_0^3 \int_{-2}^0 (x^2y - 2xy) \, dy \, dx$        $\int_{\pi}^{2\pi} \int_0^{\pi} (\sin x + \cos y) \, dx \, dy$

e-  $\iint_S (\sin x + \cos y) \, dA$       bounded by the area in fig.8

f-  $\iint_R xy^2 \, dA$       bounded by the area in fig.9

g-  $\iint_T (x - 3y) \, dA$       in fig.10

h-  $\iint_R dA$ .      bounded by the area in fig.11

i-  $\iint_R dA$ .      bounded by the area in fig.12





Figure 1




Figure 2




Figure 3




Figure 5

**Solution**

$$\begin{aligned}
 \text{a- } \int_0^3 \int_1^2 (1 + 8xy) dy dx &= \int_0^3 \left( y + 8x \frac{y^2}{2} \right) \Big|_1^2 dx \\
 &= \int_0^3 \{1 + 12x\} dx \\
 &= \left( x + 12 \frac{x^2}{2} \right) \Big|_0^3 \\
 &= (3 + 6(9)) - (0) = (3 + 54) = 57
 \end{aligned}$$

$$\int_0^1 \int_0^1 \left( 1 - \frac{x^2 + y^2}{2} \right) dx dy = \int_0^1 \left[ x - \frac{x^3}{6} - \frac{xy^2}{2} \right]_0^1 dy$$

b-

$$= \int_0^1 \left( \frac{5}{6} - \frac{y^2}{2} \right) dy = \left[ \frac{5}{6}y - \frac{y^3}{6} \right]_0^1 = \frac{2}{3}$$

$$\begin{aligned} \text{c- } \int_0^3 \int_{-2}^0 (x^2y - 2xy) dy dx &= \int_0^3 \left[ \frac{x^2y^2}{2} - xy^2 \right]_{-2}^0 dx = \int_0^3 (4x - 2x^2) dx \\ &= \left[ 2x^2 - \frac{2x^3}{3} \right]_0^3 = 0 \end{aligned}$$

$$\begin{aligned} \text{d- } \int_{\pi}^{2\pi} \int_0^{\pi} (\sin x + \cos y) dx dy &= \int_{\pi}^{2\pi} [-\cos x + x \cos y]_0^{\pi} dy : \\ &= \int_{\pi}^{2\pi} (2 + \pi \cos y) dy = [2y + \pi \sin y]_{\pi}^{2\pi} = 2\pi \end{aligned}$$

$$\begin{aligned} \text{e- } \iint_S (\sin x + \cos y) dA &= \int_0^{\pi/2} \int_0^{\pi/2} (\sin x + \cos y) dy dx \\ &= \int_0^{\pi/2} dx \left( y \sin x + \sin y \right) \Big|_{y=0}^{y=\pi/2} \\ &= \int_0^{\pi/2} \left( \frac{\pi}{2} \sin x + 1 \right) dx \\ &= \left( -\frac{\pi}{2} \cos x + x \right) \Big|_0^{\pi/2} = \frac{\pi}{2} + \frac{\pi}{2} = \pi. \end{aligned}$$

$$\begin{aligned} \text{f- } \iint_R xy^2 dA &= \int_0^1 x dx \int_{x^2}^{\sqrt{x}} y^2 dy \\ &= \int_0^1 x dx \left( \frac{1}{3} y^3 \right) \Big|_{y=x^2}^{y=\sqrt{x}} \\ &= \frac{1}{3} \int_0^1 (x^{5/2} - x^7) dx \\ &= \frac{1}{3} \left( \frac{2}{7} x^{7/2} - \frac{x^8}{8} \right) \Big|_0^1 \\ &= \frac{1}{3} \left( \frac{2}{7} - \frac{1}{8} \right) = \frac{3}{56}. \end{aligned}$$


$$\begin{aligned}
g- \iint_T (x - 3y) dA &= \int_0^a dx \int_0^{b(1-(x/a))} (x - 3y) dy \\
&= \int_0^a dx \left( xy - \frac{3}{2}y^2 \right) \Big|_{y=0}^{y=b(1-(x/a))} \\
&= \int_0^a \left[ b \left( x - \frac{x^2}{a} \right) - \frac{3}{2}b^2 \left( 1 - \frac{2x}{a} + \frac{x^2}{a^2} \right) \right] dx \\
&= \left( b \frac{x^2}{2} - \frac{b}{a} \frac{x^3}{3} - \frac{3}{2}b^2x + \frac{3}{2} \frac{b^2x^2}{a} - \frac{1}{2} \frac{b^2x^3}{a^2} \right) \Big|_0^a \\
&= \frac{a^2b}{6} - \frac{ab^2}{2}.
\end{aligned}$$

$$\begin{aligned}
h- \int_0^1 \int_{x/2}^{2x} 1 dy dx + \int_1^2 \int_{x/2}^{3-x} 1 dy dx \\
&= \int_0^1 [y]_{x/2}^{2x} dx + \int_1^2 [y]_{x/2}^{3-x} dx \\
&= \int_0^1 \left( \frac{3}{2}x \right) dx + \int_1^2 \left( 3 - \frac{3}{2}x \right) dx \\
&= \left[ \frac{3}{4}x^2 \right]_0^1 + \left[ 3x - \frac{3}{4}x^2 \right]_1^2 = \frac{3}{2}
\end{aligned}$$


$$\begin{aligned}
i- \int_0^1 \int_{-x}^{\sqrt{x}} 1 dy dx + \int_1^4 \int_{x-2}^{\sqrt{x}} 1 dy dx \\
&= \int_0^1 [y]_{-x}^{\sqrt{x}} dx + \int_1^4 [y]_{x-2}^{\sqrt{x}} dx \\
&= \int_0^1 (\sqrt{x} + x) dx + \int_1^4 (\sqrt{x} - x + 2) dx \\
&= \left[ \frac{2}{3}x^{3/2} + \frac{1}{2}x^2 \right]_0^1 + \left[ \frac{2}{3}x^{3/2} - \frac{1}{2}x^2 + 2x \right]_1^4 = \frac{13}{3}
\end{aligned}$$

### b- Poalr form

1- Find the limits of the following integral



(a)



(b)

a-  $\int_0^\pi \int_0^1 r \, dr \, d\theta$

b-  $\int_0^{\pi/4} \int_0^{\sqrt{4 \cos 2\theta}} r \, dr \, d\theta$

2- Evaluate the following

a-  $\int_{\pi/4}^{\pi/2} \int_0^{6 \csc \theta} r^2 \cos \theta \, dr \, d\theta$       b-  $\int_0^{\pi/4} \int_0^{2 \sec \theta} r^2 \sin \theta \, dr \, d\theta$

c-  $\int_{\pi/6}^{\pi/4} \int_{\csc \theta}^{\sqrt{3} \sec \theta} r \, dr \, d\theta$       d-  $\int_0^{\pi/4} \int_{\sec \theta}^{2 \cos \theta} \frac{1}{r^4} r \, dr \, d\theta$

e-  $\iint_S x \, dA$       bounded by area shown in fig.13

f-  $\iint_C y \, dA$       bounded by area shown in fig.14

g-  $\iint_A dA$       bounded by area shown in fig.15







Figure 6




Figure 8

**Solution**

$$\begin{aligned} \text{a- } \int_{\pi/4}^{\pi/2} \int_0^{6 \csc \theta} r^2 \cos \theta \, dr \, d\theta &= 72 \int_{\pi/4}^{\pi/2} \cot \theta \csc^2 \theta \, d\theta \\ &= -36 [\cot^2 \theta]_{\pi/4}^{\pi/2} = 36 \end{aligned}$$

$$\text{b- } \int_0^{\pi/4} \int_0^{2 \sec \theta} r^2 \sin \theta \, dr \, d\theta = \frac{8}{3} \int_0^{\pi/4} \tan \theta \sec^2 \theta \, d\theta = \frac{4}{3}$$

$$\begin{aligned} \text{c- } \int_{\pi/6}^{\pi/4} \int_{\csc \theta}^{\sqrt{3} \sec \theta} r \, dr \, d\theta &= \int_{\pi/6}^{\pi/4} \left( \frac{3}{2} \sec^2 \theta - \frac{1}{2} \csc^2 \theta \right) d\theta \\ &= \left[ \frac{3}{2} \tan \theta + \frac{1}{2} \cot \theta \right]_{\pi/6}^{\pi/4} = 2 - \sqrt{3} \end{aligned}$$

$$\begin{aligned} \text{d- } \int_0^{\pi/4} \int_{\sec \theta}^{2 \cos \theta} \frac{1}{r^4} r \, dr \, d\theta &= \int_0^{\pi/4} \left[ -\frac{1}{2r^2} \right]_{\sec \theta}^{2 \cos \theta} d\theta = \int_0^{\pi/4} \left( \frac{1}{2} \cos^2 \theta - \frac{1}{8} \sec^2 \theta \right) d\theta \\ &= \left[ \frac{1}{4} \theta + \frac{1}{8} \sin 2\theta - \frac{1}{8} \tan \theta \right]_0^{\pi/4} = \frac{\pi}{16} \end{aligned}$$

$$\text{e- } \int_{\sec \theta}^{\sqrt{2}} r \cos \theta \, r \, dr \, d\theta$$

$$\begin{aligned}
\iint_S x \, dA &= 2 \int_0^{\pi/4} d\theta \int_{\sec\theta}^{\sqrt{2}} r \cos\theta \, r \, dr \\
&= \frac{2}{3} \int_0^{\pi/4} \cos\theta (2\sqrt{2} - \sec^3\theta) \, d\theta \\
&= \frac{4\sqrt{2}}{3} \sin\theta \Big|_0^{\pi/4} - \frac{2}{3} \tan\theta \Big|_0^{\pi/4} \\
&= \frac{4}{3} - \frac{2}{3} = \frac{2}{3}
\end{aligned}$$

$$\begin{aligned}
\text{f- } \iint_C y \, dA &= \int_0^\pi \int_0^{1+\cos\theta} r \sin\theta \, r \, dr \, d\theta \\
&= \frac{1}{3} \int_0^\pi \sin\theta (1 + \cos\theta)^3 \, d\theta \quad \begin{array}{l} \text{Let } u = 1 + \cos\theta \\ du = -\sin\theta \, d\theta \end{array} \\
&= \frac{1}{3} \int_0^2 u^3 \, du = \frac{u^4}{12} \Big|_0^2 = \frac{4}{3}
\end{aligned}$$

$$\begin{aligned}
\text{g- } \int_{\pi/4}^{3\pi/4} \int_{\csc\theta}^{2\sin\theta} r \, dr \, d\theta &= \frac{1}{2} \int_{\pi/4}^{3\pi/4} (4\sin^2\theta - \csc^2\theta) \, d\theta \\
&= \frac{1}{2} [2\theta - \sin 2\theta + \cot\theta]_{\pi/4}^{3\pi/4} = \frac{\pi}{2}
\end{aligned}$$

### c- Change of variables

Evaluate the following integrals in polar form

$$\text{a- } \int_0^2 \int_0^{\sqrt{4-y^2}} (x^2 + y^2) \, dx \, dy$$

$$\text{b- } \int_0^2 \int_0^x y \, dy \, dx$$

$$\text{c- } \int_{\sqrt{2}}^2 \int_{\sqrt{4-y^2}}^y dx \, dy$$

d-  $\int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \frac{2}{(1+x^2+y^2)^2} dy dx$

e-  $\int_0^1 \int_x^{\sqrt{2-x^2}} (x+2y) dy dx$

f-w  $\iint_S (x+y) dA$  bounded in fig.16

g-  $\iint_T (x^2+y^2) dA$  where T is the area bounded in fig.17




Figure 10




Figure 9

**Solution**

$$a- \int_0^2 \int_0^{\sqrt{4-y^2}} (x^2 + y^2) dx dy$$


$$x_2 = \sqrt{4 - y^2} \quad , \quad x_1 = 0$$

$$y_2 = 2 \quad , \quad y_1 = 0$$

$$f(x, y) = x^2 + y^2 = r^2 \quad , \quad dxdy = r dr d\theta$$

$$\int_0^2 \int_0^{\sqrt{4-y^2}} (x^2 + y^2) dx dy = \int_{\theta_1=0}^{\theta_2=\frac{\pi}{2}} \int_{r_1=0}^{r_2=2} r^2 r dr d\theta$$

$$= \int_{\theta_1=0}^{\theta_2=\frac{\pi}{2}} \int_{r_1=0}^{r_2=2} r^3 dr d\theta = \int_0^{\pi/2} \left. \frac{r^4}{4} \right|_0^2 d\theta = 4 \int_0^{\pi/2} d\theta = 2\pi$$



$$b- \int_0^2 \int_0^x y dy dx$$


$$y_2 = x \quad , \quad y_1 = 0$$

$$x_2 = 2 \quad , \quad x_1 = 0$$

$$f(x, y) = y = r \sin \theta \quad , \quad dxdy = r dr d\theta$$

$$\int_0^2 \int_0^x y dy dx = \int_0^{\pi/4} \int_0^{2 \sec \theta} r^2 \sin \theta dr d\theta$$

$$= \frac{8}{3} \int_0^{\pi/4} \tan \theta \sec^2 \theta d\theta = \frac{4}{3}$$



$$c- \int_{\sqrt{2}}^2 \int_{\sqrt{4-y^2}}^y dx dy$$

$$x_2 = y \quad , \quad x_1 = \sqrt{4-y^2}$$


$$y_2 = 2 \quad , \quad y_1 = \sqrt{2}$$

$$f(x,y) = 1 \quad , \quad dx dy = r dr d\theta$$

$$\int_{\sqrt{2}}^2 \int_{\sqrt{4-y^2}}^y dy dx = \int_{\pi/4}^{\pi/2} \int_2^{2 \csc \theta} r dr d\theta$$

$$= \int_{\pi/6}^{\pi/4} (2 \csc^2 \theta - 2) d\theta = \left[ -2 \cot \theta - \frac{1}{2} \theta \right]_{\pi/4}^{\pi/2}$$

$$= 2 - \frac{\pi}{2}$$



$$d- \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \frac{2}{(1+x^2+y^2)^2} dy dx$$


$$y_2 = \sqrt{1-x^2} \quad , \quad y_1 = -\sqrt{1-x^2}$$

$$x_2 = 1 \quad , \quad x_1 = -1$$

$$\frac{2}{(1+x^2+y^2)^2} = \frac{2}{(1+r^2)^2} \quad , \quad dx dy = r dr d\theta$$

$$= \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \frac{2}{(1+x^2+y^2)^2} dy dx$$

$$= 4 \int_0^{\pi/2} \int_0^1 \frac{2r}{(1+r^2)^2} dr d\theta = 4 \int_0^{\pi/2} \left[ -\frac{1}{1+r^2} \right]_0^1 d\theta = 2 \int_0^{\pi/2} d\theta = \pi$$




$$e- \int_0^1 \int_x^{\sqrt{2-x^2}} (x+2y) dy dx$$

$$x_2 = 1 \quad , \quad x_1 = 0$$

$$y_2 = \sqrt{2-x^2} \quad , \quad y_1 = x$$

$$(x+2y) = r(\cos \theta + 2 \sin \theta) \quad ,$$

$$dxdy = r dr d\theta$$



$$\int_0^1 \int_x^{\sqrt{2-x^2}} (x+2y) dy dx = \int_{\pi/4}^{\pi/2} \int_0^{\sqrt{2}} (r \cos \theta + 2r \sin \theta) r dr d\theta$$

$$= \int_{\pi/4}^{\pi/2} \left[ \frac{r^3}{3} \cos \theta + \frac{2r^3}{3} \sin \theta \right]_0^{\sqrt{2}} d\theta = \int_{\pi/4}^{\pi/2} \left( \frac{2\sqrt{2}}{3} \cos \theta + \frac{4\sqrt{2}}{3} \sin \theta \right) d\theta$$

$$= \left[ \frac{2\sqrt{2}}{3} \sin \theta - \frac{4\sqrt{2}}{3} \cos \theta \right]_{\pi/4}^{\pi/2} = \frac{2(1+\sqrt{2})}{3}$$

$$f- \iint_S (x+y) dA = \int_0^{\pi/3} d\theta \int_0^a (r \cos \theta + r \sin \theta) r dr$$

$$= \int_0^{\pi/3} (\cos \theta + \sin \theta) d\theta \int_0^a r^2 dr$$

$$= \frac{a^3}{3} (\sin \theta - \cos \theta) \Big|_0^{\pi/3}$$

$$= \left[ \left( \frac{\sqrt{3}}{2} - \frac{1}{2} \right) - (-1) \right] \frac{a^3}{3} = \frac{(\sqrt{3}+1)a^3}{6}$$

$$h- \iint_T (x^2 + y^2) dA = \int_0^{\pi/4} d\theta \int_0^{\sec \theta} r^3 dr$$

$$= \frac{1}{4} \int_0^{\pi/4} \sec^4 \theta d\theta$$

$$= \frac{1}{4} \int_0^{\pi/4} (1 + \tan^2 \theta) \sec^2 \theta d\theta \quad \text{Let } u = \tan \theta \\ du = \sec^2 \theta d\theta$$

$$\begin{aligned}
 &= \frac{1}{4} \int_0^1 (1 + u^2) du \\
 &= \frac{1}{4} \left( u + \frac{u^3}{3} \right) \Big|_0^1 = \frac{1}{3}
 \end{aligned}$$

### d- Triple integral

Evaluate the following integrals:

$$a- \int_0^1 \int_0^1 \int_0^1 (x^2 + y^2 + z^2) dz dy dx$$

$$b- \int_0^{\sqrt{2}} \int_0^{3y} \int_{x^2+3y^2}^{8-x^2-y^2} dz dx dy$$

$$c-d \int_0^2 \int_{-\sqrt{4-y^2}}^{\sqrt{4-y^2}} \int_0^{2x+y} dz dx dy$$

$$\int_0^{\pi/6} \int_0^1 \int_{-2}^3 y \sin z dx dy dz$$

$$e- \int_0^1 \int_0^{1-x^2} \int_3^{4-x^2-y} x dz dy dx$$

### Solution

$$\begin{aligned}
 a- \int_0^1 \int_0^1 \int_0^1 (x^2 + y^2 + z^2) dz dy dx &= \int_0^1 \int_0^1 (x^2 + y^2 + \frac{1}{3}) dy dx \\
 &= \int_0^1 (x^2 + \frac{2}{3}) dx = 1
 \end{aligned}$$

$$\begin{aligned}
 b- \int_0^{\sqrt{2}} \int_0^{3y} \int_{x^2+3y^2}^{8-x^2-y^2} dz dx dy &= \int_0^{\sqrt{2}} \int_0^{3y} (8 - 2x^2 - 4y^2) dx dy \\
 &= \int_0^{\sqrt{2}} [8x - \frac{2}{3}x^3 - 4xy^2]_0^{3y} dy = \int_0^{\sqrt{2}} (24y - 18y^3 - 12y^3) dy \\
 &= [12y^2 - \frac{15}{2}y^4]_0^{\sqrt{2}} = 24 - 30 = -6
 \end{aligned}$$

$$\begin{aligned}
 c- \int_0^2 \int_{-\sqrt{4-y^2}}^{\sqrt{4-y^2}} \int_0^{2x+y} dz dx dy &= \int_0^2 \int_{-\sqrt{4-y^2}}^{\sqrt{4-y^2}} (2x + y) dx dy \\
 &= \int_0^2 [x^2 + xy]_{-\sqrt{4-y^2}}^{\sqrt{4-y^2}} dy = \int_0^2 (4 - y^2)^{1/2} (2y) dy
 \end{aligned}$$

$$= \left[ -\frac{2}{3} (4 - y^2)^{3/2} \right]_0^2 = \frac{2}{3} (4)^{3/2} = \frac{16}{3}$$

$$\begin{aligned} \text{d- } \int_0^{\pi/6} \int_0^1 \int_{-2}^3 y \sin z \, dx \, dy \, dz &= \int_0^{\pi/6} \int_0^1 5y \sin z \, dy \, dz \\ &= \frac{5}{2} \int_0^{\pi/6} \sin z \, dz = \frac{5(2 - \sqrt{3})}{4} \end{aligned}$$

$$\begin{aligned} \text{e- } \int_0^1 \int_0^{1-x^2} \int_3^{4-x^2-y} x \, dz \, dy \, dx &= \int_0^1 \int_0^{1-x^2} x(1 - x^2 - y) \, dy \, dx \\ &= \int_0^1 x \left[ (1 - x^2)^2 - \frac{1}{2} (1 - x^2) \right] dx = \int_0^1 \frac{1}{2} x (1 - x^2)^2 dx \\ &= \left[ -\frac{1}{12} (1 - x^2)^3 \right]_0^1 = \frac{1}{12} \end{aligned}$$

### e- Surface area

Find the area of the following surfaces:

a-  $Z = f(x, y) = 6 - 3x - 2y$  lies in the region shown in fig. 18

b-  $Z = x^2 + y^2$  lies in the region shown in fig. 19




Figure 11




Figure 12

### Solution

a-  $Z = 6 - 3x - 2y$

$$\frac{\partial f}{\partial x} = -3 \quad , \quad \frac{\partial f}{\partial y} = -2 \quad , \quad 0 \leq x \leq 2 \quad , \quad 0 \leq y \leq -\frac{3}{2}x + 3$$



$$\begin{aligned}
 S &= \iint_R \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dA \\
 &= \int_0^2 \int_0^{-\frac{3}{2}x+3} \sqrt{(-3)^2 + (-2)^2 + 1} dy dx = \sqrt{14} \int_0^2 \left(-\frac{3}{2}x + 3\right) dx \\
 S &= \sqrt{14} \left(-\frac{3}{4}x^2 + 3x\right) \Big|_0^2 = 3\sqrt{14}
 \end{aligned}$$

b-  $Z = x^2 + y^2$

$$\frac{\partial f}{\partial x} = 2x \quad , \quad \frac{\partial f}{\partial y} = 2y$$

$$\begin{aligned}
 S &= \iint_R \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dA \\
 &= \iint_R \sqrt{1 + 4x^2 + 4y^2} dA \quad r^2 = x^2 + y^2 \\
 &= \int_0^{2\pi} \int_0^3 r \sqrt{1 + 4r^2} dr d\theta = \frac{2}{24} \int_0^{2\pi} [1 + 4r^2]^{\frac{3}{2}} \Big|_0^3 d\theta = 18 \int_0^{2\pi} d\theta = 36\pi
 \end{aligned}$$