



Estimation of Creatinine in serum

Creatinine is a chemical waste product that's produced by your muscle metabolism and to a smaller extent by eating meat. Healthy kidneys filter creatinine and other waste products from your blood. The filtered waste products leave your body in your urine.

The creatinine blood test helps doctors to diagnose kidney disease. A poorly functioning kidney cannot filter creatinine as well as it usually does, which causes levels in the blood to rise.

* The result of this blood test is useful, as it is an important marker of how well the kidneys are working.

The production of creatinine depends on muscle mass and the amount of creatinine increases in muscle size and also depends on the extent of glomerular filtration in the kidney.

Causes For High Levels

Some of the causes of high creatinine levels are:

1. Chronic kidney disease

When kidneys are damaged, they have trouble removing creatinine from the blood and levels rise.

2. Kidney obstruction

A blockage in the flow of urine, such as an enlarged prostate or kidney stone, could cause kidney obstruction which might raise the level of creatinine. The medical term for this condition is hydronephrosis.



3. Dehydration

Severe dehydration is a risk factor for kidney injury, which will affect creatinine levels.

4. Increased consumption of protein

What a person eats can have a significant impact on creatinine levels. For example, proteins and cooked meat contain creatinine, so eating more than the recommended amount of meat or other proteins for your activity levels, or adding extra protein to the diet through supplements can cause high creatinine levels.

5. Intense exercise

Creatine is present in the muscles and helps them produce energy. Rigorous exercise can increase creatinine levels by increasing muscle breakdown.

6. Certain medications

Antibiotics, such as trimethoprim, and H₂ blockers, such as cimetidine, can cause a temporary increase in measured serum creatinine levels.

Causes For Low Levels

1. Low muscle mass

Because the breakdown of muscle produces creatinine, low muscle mass can result in low levels of creatinine.

Older people are more at risk as muscle mass declines with age. Malnutrition can also cause low muscle mass and low creatinine levels.



2. Pregnancy

Pregnancy causes an increase in blood flow to the kidney leading to increased urine production and faster elimination of creatinine, leading to lower levels.

3. Extreme weight loss

Weight loss can result in the reduction of muscle mass, leading to low levels of creatinine.

Normal level:

Men: The proportion of natural creatine have higher than in women, because they have a larger muscle mass and stronger, so the proportion of creatinine have a natural range of 0.6-1.2 mg\dl, in adult males.

For women: In adult women it ranges from 0.5-1.1 mg\dl

Vegetarians: They have a low creatinine ratio compared to people who eat meat.

Men with muscle: the ratio is generally higher.

Older people: The percentage is generally lower.

Infants: Their normal ratio starts at 0.2 mg\dl and starts to increase as their muscles develop.

People with one kidney: the normal ratio is 1.8 or 1.9 mg\dl