



## Review

\***Molecular Biology** is the field of biology that studies the composition, structure and interactions of cellular molecules – such as nucleic acids and proteins – that carry out the biological processes essential for the cell's functions and maintenance.

\***Nucleic acid** is an important class of macromolecules found in all cells and viruses. The functions of nucleic acids have to do with the **storage and expression of genetic information**.

\*There are two types of nucleic acids, deoxyribonucleic acid (**DNA**) and ribonucleic acid (**RNA**).

\* Nucleic acids are the polymers of nucleotides( built up by the monomeric units called **nucleotides** - polynucleotides).


\*Nucleotide essentially consists of **base, sugar** and **phosphate**. The term nucleoside refers to **base + sugar**. Thus, nucleotide is **nucleoside + phosphate**.

\*The nitrogenous bases found in nucleotides are aromatic heterocyclic compounds. The bases are of two types **purines** (Adenine and Guanine) and **pyrimidine's**(Thymine , Cytosine and Uracil) .

- ADENINE pairs with THYMINE (A::T) with two hydrogen bonds
- GUANINE pairs with CYTOSINE (G::C) with three hydrogen bonds
- DNA is composed of four nucleotides, each containing: adenine, cytosine, thymine, or guanine.


- The amounts of A = T, G = C, and purines = pyrimidine [Chargaff's Rule].
- DNA is a double-stranded helix with antiparallel strands [Watson and Crick].
- Nucleotides in each strand are linked by 5'-3' phosphodiester bonds.
- Bases on opposite strands are linked by hydrogen bonding: A with T, and G with C.

\*The interrelationship of these three classes of biomolecules (DNA, RNA and proteins) constitutes the central dogma of molecular biology or more commonly the central dogma of life.



## DNA Replication


- DNA replication, the basis for biological inheritance, is a fundamental process occurring in all living organisms to copy their DNA.
- In the process of "replication" each strand of the original double-stranded DNA molecule serves as template for the reproduction of the complementary strand.
- Two identical DNA molecules have been produced from a single double-stranded DNA molecule.
- In a cell, DNA replication begins at specific locations in the genome, called "origins".
- \* A eukaryotic chromosome may have hundreds or even thousands of replication origins
- Unwinding of DNA at the origin, and synthesis of new strands, forms a replication fork.
- In addition to DNA polymerase, the enzyme that synthesizes the new DNA by adding nucleotides matched to the template strand, a number of other proteins are associated with the fork and assist in the initiation and continuation of DNA synthesis.





## The Mechanism of DNA Replication

- DNA synthesis on the leading strand is continuous.
- The lagging strand grows the same general direction as the leading strand (in the same direction as the Replication Fork). However, DNA is made in the 5'-to-3' direction.
- Therefore, DNA synthesis on the lagging strand is discontinuous
- DNA is added as short fragments (Okasaki fragments) that are subsequently ligated together.





## Enzymes in DNA replication

Many proteins assist in DNA replication

- DNA **helicases** unwind the double helix, the template strands are stabilized by other Proteins
- Single-stranded DNA binding proteins make the template available
- RNA **primase** catalyzes the synthesis of short RNA primers, to which nucleotides are added.
- DNA **polymerase III** extends the strand in the 5'-to-3' direction.
- DNA **polymerase I** degrades the RNA primer and replaces it with DNA.
- DNA **ligase** joins the DNA fragments into a continuous daughter strand.