
Mathematics (II)

Dr. Alaa Mohammed Hussein Wais

## **13. Tangent planes and normal lines:-**



The tangent plane at the point  $P_o(x_o, y_o, z_o)$  on the level surface f(x, y, z) = C of a differentiable function f is the plane through  $P_o$  normal to  $\nabla f|_P$ 

The normal line of the surface at  $P_o$  is the line through  $P_o$  parallel to  $\nabla f|_P$ 

The tangent plane and normal line have the following equation:

**Tangent plane** to f(x, y, z) = C at  $P_o(x_o, y_o, z_o)$ :  $f_x(p_o)(x - x_o) + f_y(p_o)(y - y_o) + f_z(p_o)(z - z_o) = 0$ 

Normal line to f(x, y, z) = C at  $P_o(x_o, y_o, z_o)$ :  $x = x_o + f_x(p_o)t$ ,  $y = y_o + f_y(p_o)t$ ,  $z = z_o + f_z(p_o)t$  Mathematics (II ) Al-mustaqbal University Collage Dr. *Alaa Mohammed Hussein Wais* 

*Example*: find the tangent plane and normal line of the surface  $f(x, y, z) = x^2 + y^2 + z - 9 = 0$  at the point  $P_o(1, 2, 4)$ 

Solution: the tangent plane is:

 $f_x(p_o)(x-x_o) + f_y(p_o)(y-y_o) + f_z(p_o)(z-z_o) = 0$ 

$$f_x = \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} (x^2 + y^2 + z - 9) = 2x$$
  

$$f_x(P_o) = f_x(1,2,4) = (2)(1) = 2$$
  

$$f_y = \frac{\partial f}{\partial y} = \frac{\partial}{\partial y} (x^2 + y^2 + z - 9) = 2y$$
  


$$f_y(P_o) = f_y(1,2,4) = (2)(2) = 4$$
  

$$f_z = \frac{\partial f}{\partial z} = \frac{\partial}{\partial z} (x^2 + y^2 + z - 9) = 1$$
  

$$f_z(P_o) = f_z(1,2,4) = 1$$

Mathematics (II)

Dr. Alaa Mohammed Hussein Wais



*Example*: find the tangent plane and normal line of the surface  $f(x, y, z) = x^2 + y^2 + z^2 = 3$  at the point  $P_o(1,1,1)$ 

Solution: the tangent plane is:

$$f_x(p_o)(x - x_o) + f_y(p_o)(y - y_o) + f_z(p_o)(z - z_o) = 0$$

$$\begin{split} f_x &= \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} (x^2 + y^2 + z^2) = 2x \\ f_x(P_o) &= f_x(1,1,1) = (2)(1) = \boxed{2} \\ f_y &= \frac{\partial f}{\partial y} = \frac{\partial}{\partial y} (x^2 + y^2 + z^2) = 2y \\ f_y(P_o) &= f_y(1,1,1) = (2)(1) = \boxed{2} \\ f_z &= \frac{\partial f}{\partial z} = \frac{\partial}{\partial z} (x^2 + y^2 + z^2) = 2z \\ f_z(P_o) &= f_z(1,1,1) = (2)(1) = \boxed{2} \end{split}$$

Mathematics (II)

Dr. Alaa Mohammed Hussein Wais

.. The tangent plane is:  

$$2(x-1)+2(y-1)+2(z-1)=0$$
or  

$$2x-2+2y-2+2z-2=0$$

$$2x+2y+2z-6=0$$

$$2x+2y+2z=6$$

$$2(x+y+z)=6 \implies x+y+z=3$$
The normal line is:

$$x = x_o + f_x(p_o)t , y = y_o + f_y(p_o)t , z = z_o + f_z(p_o)t$$
  

$$\therefore x = 1 + 2t , y = 1 + 2t , z = 1 + 2t$$

*H.W*: find the tangent plane and normal line of the surface  $f(x, y, z) = x^2 + 2xy - y^2 + z^2 = 7$  at the point  $P_o(1, -1, 3)$