




## Directional derivatives :

Suppose that the function  $f(x, y)$  is defined throughout a region  $R$  in the  $xy$  - plane, that  $P_o(x_o, y_o)$  is a point in  $R$  and that  $u = u_1\mathbf{i} + u_2\mathbf{j}$  is a unit vector. Thus the equations:

$$x = x_o + su_1 \quad , \quad y = y_o + su_2$$

Parameterize the line through  $P_o$  parallel to  $u$ . if the parameter  $s$  measures arc length from  $P_o$  in the direction of  $u$ , we find the rate of change of  $f$

at  $P_o$  in the direction of  $u$  by calculating  $\frac{df}{ds}$  at  $P_o$  at



## *Gradient vector:*

$$f(x, y, z)$$

$$P_o(x_o, y_o, z_o)$$

$$\nabla f = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j} + \frac{\partial f}{\partial z} \mathbf{k}$$



The notation  $\nabla f$  is read ( grad  $f$ ) as well as ( gradient  $f$ ) and ( del  $f$ ).  
The symbol  $\nabla$  by itself is read (del)

### *The directional derivatives*

If  $f(x, y, z)$  has continuous partial derivatives at  $P_o(x_o, y_o, z_o)$  and  $u$  is a unit vector, then the derivative of  $f$  at  $P_o$  in the direction of  $u$  is:

$$(DuF)|_{p_o} = \nabla f|_{p_o} \cdot u$$

Which is the scalar product of the gradient of  $F$  at  $P_o$  and  $u$

" $(DuF)|_{p_o} \rightarrow$  The derivative of  $F$  at  $P_o$  in the direction of  $u$ "

**Example:** Find the direction derivative of the function

$F(x, y, z) = x^2 + y^2 + z^2$  at point  $p_o(1, 1, 1)$  in the direction of

Vector  $v = i + j + k$ .

**Solution //**

$$(DuF)|_{p_o} = \nabla f|_{p_o} \cdot u$$

$$\nabla f = \frac{\partial f}{\partial x} i + \frac{\partial f}{\partial y} j + \frac{\partial f}{\partial z} k = 2x i + 2y j + 2z k$$

$$\text{at point } p_o(1, 1, 1) \rightarrow \nabla f|_{p_o} = 2i + 2j + 2k$$

$$u = \frac{v}{|v|} = \frac{i+j+k}{\sqrt{1+1+1}} = \frac{i+j+k}{\sqrt{3}}$$

$$(DuF)|_{p_o} = \nabla f|_{p_o} \cdot u = 2i + 2j + 2k \cdot \frac{i+j+k}{\sqrt{3}} = \frac{6}{\sqrt{3}}$$



**Example:** find the derivative of  $f(x, y) = xe^y + \cos(xy)$  at the point  $(2, 0)$  in the direction of  $v = 3i - 4j$

**Solution:** the direction of  $v$  is the unit vector obtained by dividing  $v$  by its length:

$$u = \frac{v}{|v|}$$

$$|v| = \sqrt{(3)^2 + (-4)^2} = \sqrt{9+16} = \sqrt{25} = 5$$

$$u = \frac{3i - 4j}{5} = \frac{3}{5}i - \frac{4}{5}j$$

$$\nabla f = \frac{\partial f}{\partial x}i + \frac{\partial f}{\partial y}j = (e^y - y \sin xy)i + (xe^y - x \sin xy)j$$

$$\text{at point } p_0(2, 0) \rightarrow \nabla f|_{p_0} = i + 2j$$

$$(DuF)|_{p_0} = \nabla f|_{p_0} \cdot u = i + 2j \cdot \frac{3i-4j}{5} = \frac{-5}{5} = -1$$

**Example:** find the derivative of function  $f(x, y) = 2xy - 3y^2$  at the point  $P_0(5, 5)$  in the direction of  $A = 4i + 3j$

**Solution:**

$$u = \frac{A}{|A|}$$

$$|A| = \sqrt{(4)^2 + (3)^2} = \sqrt{16+9} = \sqrt{25} = 5$$

$$u = \frac{4i+3j}{5} = \frac{4}{5}i + \frac{3}{5}j$$

$$\nabla f = \frac{\partial f}{\partial x}i + \frac{\partial f}{\partial y}j = (2y)i + (2x - 6y)j$$

$$\text{at point } p_0(5, 5) \rightarrow \nabla f|_{p_0} = 10i - 20j$$

$$(DuF)|_{p_0} = \nabla f|_{p_0} \cdot u = 10i - 20j \cdot \frac{4i+3j}{5} = \frac{-20}{5} = -4$$



**H.W:**

1. find the derivative of the function  $f(x, y, z) = xy + yz + zx$ , at the point  $P_0(1, -1, 2)$  in the direction of  $A = 3i + 6j - 2k$
  
2. find the derivative of the function  $g(x, y, z) = 3e^x \cos yz$ , at the point  $P_0(0, 0, 0)$  in the direction of  $A = 2i + j - 2k$